

ABSTRACT AND REFERENCES

TECHNOLOGY AND EQUIPMENT OF FOOD PRODUCTION

DOI: 10.15587/1729-4061.2019.160021

CONSTRUCTION OF A METHOD FOR PREDICTING THE NUMBER OF ENTEROBACTERIA IN MILK USING ARTIFICIAL NEURAL NETWORKS (p. 6-13)**Oleksandra Berhilevych**

Sumy State University, Sumy, Ukraine

ORCID: <http://orcid.org/0000-0002-3622-8942>**Victoria Kasianchuk**

Sumy State University, Sumy, Ukraine

ORCID: <http://orcid.org/0000-0001-5849-7679>**Ihor Chernetskyi**

Sumy State University, Sumy, Ukraine

ORCID: <http://orcid.org/0000-0002-7737-8437>**Anastasia Konieva**

Sumy State University, Sumy, Ukraine

ORCID: <http://orcid.org/0000-0001-8516-1353>**Lubov Dimitrijevič**

Sumy National Agrarian University, Sumy, Ukraine

ORCID: <http://orcid.org/0000-0003-3616-1167>**Tatyana Marenkova**

Sumy National Agrarian University, Sumy, Ukraine

ORCID: <http://orcid.org/0000-0001-7481-0848>

It is now established that artificial neural networks (ANNs) provide better simulation and prediction of the number of microorganisms in raw materials and foodstuffs. In this case, ANNs could be used as informative, fast, and cost-effective means. According to the European requirements to food products, basic microbiological indicators are the total number of microorganisms and bacteria from the Enterobacteriaceae family, since they are most commonly associated with food-borne diseases and poisonings. The aim of this work was to devise a method for predicting the number of bacteria from the Enterobacteriaceae family in raw milk at its chilled storage and to estimate the predictive capability of ANN. Construction of the method included 4 stages. At the first stage, we examined the number of enterobacteria depending on the physical-chemical composition of raw milk, temperature and duration of storage in a refrigerator. At the second stage, we compiled a base of experimental data obtained from research models. At the next stage, we introduced the received database to ANN. And at the last stage we assessed effectiveness of the predicting technique. The constructed ANN consists of three layers: an input layer (5 parameters: milk storage temperature (4, 6, 8, and 10 °C), duration of milk storage (from 1 to 48 hours); the acidity of milk (17–20 %), the fat content (3.2; 3.6; 4.0; 4.5 %) and protein content (2.9; 3.0; 3.3 %) in milk; hidden layers (with 30 neurons) and the output layer (the projected number of bacteria). In order to train and optimize the ANN, we used 1,200 experimental data, which revealed that the prediction had the highest rate of deviation of 2.497 % (or 370 bacterial cells per 1 ml). Thus, the devised predicting method could be used to predict the number of bacteria taking into consideration the complex of environmental variables in different food products. In addition, a given approach could be employed as artificial intelligence when assessing microbiological risks and for quick monitoring of food safety.

Keywords: Enterobacteriaceae, raw milk, artificial neural networks, predicting the number of bacteria.

References

- Costard, S., Espejo, L., Groenendaal, H., Zagmutt, F. J. (2017). Outbreak-Related Disease Burden Associated with Consumption of Unpasteurized Cow's Milk and Cheese, United States, 2009–2014. *Emerging Infectious Diseases*, 23(6), 957–964. doi: <https://doi.org/10.3201/eid2306.151603>
- Fox, E. M., Jiang, Y., Gobius, K. S. (2018). Key pathogenic bacteria associated with dairy foods: On-farm ecology and products associated with foodborne pathogen transmission. *International Dairy Journal*, 84, 28–35. doi: <https://doi.org/10.1016/j.idairyj.2018.03.013>
- Van der Meulen, B. (2013). The Structure of European Food Law. *Laws*, 2 (2), 69–98. doi: <https://doi.org/10.3390/laws2020069>
- Rahmat, S., Cheong, C. B., Hamid, M. S. R. B. A. (2016). Challenges of Developing Countries in Complying Quality and Enhancing Standards in Food Industries. *Procedia – Social and Behavioral Sciences*, 224, 445–451. doi: <https://doi.org/10.1016/j.sbspro.2016.05.418>
- Faour-Klingbeil, D., Todd, E. (2018). A Review on the Rising Prevalence of International Standards: Threats or Opportunities for the Agri-Food Produce Sector in Developing Countries, with a Focus on Examples from the MENA Region. *Foods*, 7 (3), 33. doi: <https://doi.org/10.3390/foods7030033>
- McMeekin, T., Olley, J., Ratkowsky, D., Corkrey, R., Ross, T. (2013). Predictive microbiology theory and application: Is it all about rates? *Food Control*, 29 (2), 290–299. doi: <https://doi.org/10.1016/j.foodcont.2012.06.001>
- Teleken, J. T., Robazza, W. da S., Gomes, G. de A. (2011). Mathematical modeling of microbial growth in milk. *Food Science and Technology (Campinas)*, 31 (4), 891–896. doi: <https://doi.org/10.1590/s0101-20612011000400010>
- Huang, L. (2014). IPMP 2013 – A comprehensive data analysis tool for predictive microbiology. *International Journal of Food Microbiology*, 171, 100–107. doi: <https://doi.org/10.1016/j.ijfoodmicro.2013.11.019>
- Pal, M., Mulu, S., Tekle, M., Pintoo, S. V., Prajapati, J. P. (2016). Bacterial contamination of dairy products. *Beverage & Food World*, 43 (9), 40–43.
- Berhilevych, O., Kasianchuk, V. (2017). Identification of Cronobacter spp (enterobacter sakazakii) from raw milk and environmental samples of dairy farms. *Eastern-European Journal of Enterprise Technologies*, 6 (11 (90)), 4–10. doi: <https://doi.org/10.15587/1729-4061.2017.114637>
- Kukhtyn, M., Berhilevych, O., Kravcheniuk, K., Shynkaruk, O., Horyuk, Y., Semaniuk, N. (2017). Formation of biofilms on dairy equipment and the influence of disinfectants on them. *Eastern-European Journal of Enterprise Technologies*, 5 (11 (89)), 26–33. doi: <https://doi.org/10.15587/1729-4061.2017.110488>
- Ray, B., Bhunia, A. (2013). *Fundamental Food Microbiology*. CRC Press, 663. doi: <https://doi.org/10.1201/b16078>
- Von Neubeck, M., Baur, C., Krewinkel, M., Stoeckel, M., Kranz, B., Stressler, T. et. al. (2015). Biodiversity of refrigerated raw milk microbiota and their enzymatic spoilage potential. *International Journal of Food Microbiology*, 211, 57–65. doi: <https://doi.org/10.1016/j.ijfoodmicro.2015.07.001>
- Zastempowska, E., Grajewski, J., Twaru ek, M. (2016). Food-Borne Pathogens and Contaminants in Raw Milk – A Review. *Annals of Animal Science*, 16 (3), 623–639. doi: <https://doi.org/10.1515/aoas-2015-0089>
- Melini, F., Melini, V., Luziatelli, F., Ruzzi, M. (2017). Raw and Heat-Treated Milk: From Public Health Risks to Nutritional Quality. *Beverages*, 3 (4), 54. doi: <https://doi.org/10.3390/beverages3040054>
- Wanjala, W. N., Nduko, J. M., Mwendu, M. C. (2018). Coliforms Contamination and Hygienic Status of Milk Chain in Emerging Economies. *Journal of Food Quality and Hazards Control*, 5 (1), 3–10. doi: <https://doi.org/10.29252/jfqc.5.1.3>

17. Paludetti, L. F., Jordan, K., Kelly, A. L., Gleeson, D. (2018). Evaluating the effect of storage conditions on milk microbiological quality and composition. *Irish Journal of Agricultural and Food Research*, 57 (1), 52–62. doi: <https://doi.org/10.1515/ijaf-2018-0006>
18. Panigrahi, S., Devi, B., Swain, K., Priyadarshini, P. (2018). Microbiology of milk: Public health aspect. *The Pharma Innovation Journal*, 7 (1), 260–264.
19. O'Connell, A., Ruegg, P. L., Jordan, K., O'Brien, B., Gleeson, D. (2016). The effect of storage temperature and duration on the microbial quality of bulk tank milk. *Journal of Dairy Science*, 99 (5), 3367–3374. doi: <https://doi.org/10.3168/jds.2015-10495>
20. Christiansson, A. (2017). A predictive model to evaluate the impact of the cooling profile on growth of psychrotrophic bacteria in raw milk from conventional and robotic milking. *Journal of Dairy Research*, 84 (03), 318–321. doi: <https://doi.org/10.1017/s0022029917000334>
21. De Souza Sant'Ana, A. (Ed.) (2017). *Quantitative microbiology in food processing: modeling the microbial ecology*. John Wiley & Sons, Ltd., 696. doi: <https://doi.org/10.1002/9781118823071>
22. Bevilacqua, A., Speranza, B., Sinigaglia, M., Corbo, M. (2015). A Focus on the Death Kinetics in Predictive Microbiology: Benefits and Limits of the Most Important Models and Some Tools Dealing with Their Application in Foods. *Foods*, 4 (4), 565–580. doi: <https://doi.org/10.3390/foods4040565>
23. Koseki, S. (2016). Predictive Modeling for Estimation of Bacterial Behavior from Farm to Table. *Food Safety*, 4 (2), 33–44. doi: <https://doi.org/10.14252/foodsafetyfscj.2016006>
24. Funes, E., Allouche, Y., Beltrán, G., Jiménez, A. (2015). A Review: Artificial Neural Networks as Tool for Control Food Industry Process. *Journal of Sensor Technology*, 05 (01), 28–43. doi: <https://doi.org/10.4236/jst.2015.51004>
25. Prieto, A., Prieto, B., Ortigosa, E. M., Ros, E., Pelayo, F., Ortega, J., Rojas, I. (2016). Neural networks: An overview of early research, current frameworks and new challenges. *Neurocomputing*, 214, 242–268. doi: <https://doi.org/10.1016/j.neucom.2016.06.014>
26. Correa, D. A., Montero Castillo, P. M., Martelo, R. J. (2018). Neural networks in food industry. *Contemporary Engineering Sciences*, 11 (37), 1807–1826. doi: <https://doi.org/10.12988/ces.2018.84141>
27. Orawan, C., Panwadee, S., Bandit, S. (2016). Application of artificial neural networks on growth prediction of *Staphylococcus aureus* in milk. *International Food Research Journal*, 23 (1), 415–418.
28. Østergaard, N. B., Dalgaard, P., Christiansen, L. E. (2014). *Predictive Food Microbiology: new tools for risk assessment and dairy product development*. DTU Food, 206.
29. Berhilevych, O. M., Kasyanchuk, V. V., Terekhyna, O. V., Berhilevych, O. O., Cherneckii, I. (2016). Prediction of number of psychrotrophic microorganisms in refrigerated cow's milk with using artificial neural networks. *Veterynarna biotekhnolohiya*, 29, 43–54.
30. Bogdanovičová, K., Vyletětlová-Klimešová, M., Babák, V., Kalhotka, L., Koláčková, I., Karpíšková, R. (2016). Microbiological Quality of Raw Milk in the Czech Republic. *Czech Journal of Food Sciences*, 34 (3), 189–196. Available at: <https://agriculturejournals.cz/publicFiles/185247.pdf>
31. Meena, G. S., Majumdar, G. C., Banerjee, R., Kumar, N., Meena, P. K. (2014). Growth Characteristics Modeling of Mixed Culture of *Bifidobacterium bifidum* and *Lactobacillus acidophilus* using Response Surface Methodology and Artificial Neural Network. *Brazilian Archives of Biology and Technology*, 57 (6), 962–970. doi: <https://doi.org/10.1590/s1516-8913201402657>
32. Sarkar, S. (2016). Microbiological Safety Concerns of Raw Milk. *Journal of Food Nutrition and Dietetics*, 01 (02), 105. doi: <https://doi.org/10.19104/jfnd.2016.105>
33. Martin, N. H., Trnčič, A., Hsieh, T.-H., Boor, K. J., Wiedmann, M. (2016). The Evolving Role of Coliforms As Indicators of Unhygienic

Processing Conditions in Dairy Foods. *Frontiers in Microbiology*, 7. doi: <https://doi.org/10.3389/fmicb.2016.01549>

34. Jaganathan, A., Kuppuraj, S. (2016). Artificial Neural Networks Methods for Predicting the Performance and Process in the Milk Industry. *International Research Journal of Engineering and Technology*, 3 (11), 1127–1132.

DOI: 10.15587/1729-4061.2019.160142

MODELING THE PROCESS OF MICROBIAL BIOFILM FORMATION ON STAINLESS STEEL WITH A DIFFERENT SURFACE ROUGHNESS (p. 14-21)

Mykola Kukhtyn

Ternopil Ivan Puluj National Technical University, Ternopil, Ukraine
ORCID: <http://orcid.org/0000-0002-0195-0767>

Khrystyna Kravcheniuk

Ternopil Ivan Puluj National Technical University, Ternopil, Ukraine
ORCID: <http://orcid.org/0000-0002-7547-6834>

Ludmila Beyko

Ternopil Ivan Puluj National Technical University, Ternopil, Ukraine
ORCID: <http://orcid.org/0000-0001-6211-8010>

Yulia Horiuk

State Agrarian and Engineering University in Podilya,
Kamianets-Podilskyi, Ukraine
ORCID: <http://orcid.org/0000-0002-7162-8992>

Oleksandr Skliar

Sumy National Agrarian University, Sumy, Ukraine
ORCID: <http://orcid.org/0000-0002-0111-1277>

Serhii Kernychnyi

State Agrarian and Engineering University in Podilya,
Kamianets-Podilskyi, Ukraine
ORCID: <http://orcid.org/0000-0001-9533-684X>

This paper reports a study into the process of bacteria adhesion to surfaces of different roughness depending on sizes and shapes. It was determined that at the surface of stainless steel with a roughness of $2.687 \pm 0.014 \mu\text{m}$, the film formation process in *E. coli* and *S. aureus* occurred equally over 3 to 24 hours and did not depend on the size of bacteria. This makes it possible to argue that rod-shaped and coccoid bacteria freely attach themselves in the grooves of roughness, followed by the initial process related to the first stage of a biofilm formation. During sanitization, the hollows of roughness could host both the coccoid and rod-shaped bacteria. At the surface of steel with a roughness of $0.95 \pm 0.092 \mu\text{m}$ the process of film formation in *S. aureus* occurred more intensely than in *E. coli*. Within 3 h of incubation, density of the formed biofilms *S. aureus* was 1.2 times larger compared to the biofilms *E. coli*. Over the following 15 hours of incubation, the biofilms *S. aureus* were on average 1.3 times denser. This suggests that *S. aureus*, due to a spherical form, can stay put in the hollows of roughness of $0.95 \pm 0.092 \mu\text{m}$ and quicker attach to the surface. At the same time, *E. coli*, owing to a rod-shaped shape, would attach at such a surface roughness to the hollows lengthwise only. It has been proven that at a surface roughness of $0.63 \pm 0.087 \mu\text{m}$ the intensity of the *S. aureus* film formation was on average 1.4 times faster than in *E. coli*. However, at a roughness of $0.16 \pm 0.018 \mu\text{m}$ the process of film formation occurred equally in *E. coli* and *S. aureus*, but the biofilms demonstrated lower density compared to those that formed at a roughness of $0.63 \pm 0.087 \mu\text{m}$.

Thus, the use of equipment with a roughness of less than $0.5 \mu\text{m}$ in the dairy industry will make it possible to reduce the attachment of microorganisms to surface and to decrease contamination of dairy products.

Keywords: microbial adhesion, formation of biofilms, roughness of stainless-steel surface, a film formation process.

References

- Kukhtyn, M., Berhilevych, O., Kravcheniuk, K., Shynkaruk, O., Horyuk, Y., Semaniuk, N. (2017). Formation of biofilms on dairy equipment and the influence of disinfectants on them. *Eastern-European Journal of Enterprise Technologies*, 5 (11 (89)), 26–33. doi: <https://doi.org/10.15587/1729-4061.2017.110488>
- Kukhtyn, M., Berhilevych, O., Kravcheniuk, K., Shynkaruk, O., Horyuk, Y., Semaniuk, N. (2017). The influence of disinfectants on microbial biofilms of dairy equipment. *EUREKA: Life Sciences*, 5, 11–17. doi: <https://doi.org/10.21303/2504-5695.2017.00423>
- Shaheen, R., Svensson, B., Andersson, M. A., Christiansson, A., Salkinoja-Salonen, M. (2010). Persistence strategies of *Bacillus cereus* spores isolated from dairy silo tanks. *Food Microbiology*, 27 (3), 347–355. doi: <https://doi.org/10.1016/j.fm.2009.11.004>
- Council directive 93/43/EEC on the hygiene of foodstuffs (1993). *Official Journal of the European Communities*, No L 175/1.
- Whitehead, K. A., Verran, J. (2007). The effect of surface properties and application method on the retention of *Pseudomonas aeruginosa* on uncoated and titanium-coated stainless steel. *International Biodeterioration & Biodegradation*, 60 (2), 74–80. doi: <https://doi.org/10.1016/j.ibiod.2006.11.009>
- Verran, J., Packer, A., Kelly, P., Whitehead, K. A. (2010). The retention of bacteria on hygienic surfaces presenting scratches of microbial dimensions. *Letters in Applied Microbiology*, 50 (3), 258–263. doi: <https://doi.org/10.1111/j.1472-765x.2009.02784.x>
- Rajab, F. H., Liauw, C. M., Benson, P. S., Li, L., Whitehead, K. A. (2018). Picosecond laser treatment production of hierarchical structured stainless steel to reduce bacterial fouling. *Food and Bioproducts Processing*, 109, 29–40. doi: <https://doi.org/10.1016/j.fbp.2018.02.009>
- Finkel, J. S., Mitchell, A. P. (2010). Genetic control of *Candida albicans* biofilm development. *Nature Reviews Microbiology*, 9 (2), 109–118. doi: <https://doi.org/10.1038/nrmicro2475>
- Monds, R. D., O'Toole, G. A. (2009). The developmental model of microbial biofilms: ten years of a paradigm up for review. *Trends in Microbiology*, 17 (2), 73–87. doi: <https://doi.org/10.1016/j.tim.2008.11.001>
- Zhao, K., Tseng, B. S., Beckerman, B., Jin, F., Gibiansky, M. L., Harrison, J. J. et al. (2013). Psl trails guide exploration and microcolony formation in *Pseudomonas aeruginosa* biofilms. *Nature*, 497 (7449), 388–391. doi: <https://doi.org/10.1038/nature12155>
- Oliveira, N. M., Martinez-Garcia, E., Xavier, J., Durham, W. M., Kolter, R., Kim, W., Foster, K. R. (2015). Correction: Biofilm Formation As a Response to Ecological Competition. *PLOS Biology*, 13 (8), e1002232. doi: <https://doi.org/10.1371/journal.pbio.1002232>
- Monds, R. D., O'Toole, G. A. (2009). The developmental model of microbial biofilms: ten years of a paradigm up for review. *Trends in Microbiology*, 17 (2), 73–87. doi: <https://doi.org/10.1016/j.tim.2008.11.001>
- Moriarty, T. F., Poulsson, A. H. C., Rochford, E. T. J., Richards, R. G. (2011). Bacterial Adhesion and Biomaterial Surfaces. *Comprehensive Biomaterials*, 75–100. doi: <https://doi.org/10.1016/b978-0-08-055294-1.00007-6>
- Hoevar, M., Jenko, M., Godec, M., Drobne, D. (2014). An overview of the influence of stainless-steel surface properties on bacterial adhesion. *Materials and technology*, 48 (5), 609–617.
- Crawford, R. J., Webb, H. K., Truong, V. K., Hasan, J., Ivanova, E. P. (2012). Surface topographical factors influencing bacterial attachment. *Advances in Colloid and Interface Science*, 179–182, 142–149. doi: <https://doi.org/10.1016/j.cis.2012.06.015>
- Merritt, K., An, Y. H. (2000). Factors Influencing Bacterial Adhesion. *Handbook of Bacterial Adhesion*, 53–72. doi: <https://doi.org/10.1385/1-59259-224-4:53>
- Garcia, S., Trueba, A., Vega, L. M., Madariaga, E. (2016). Impact of the surface roughness of AISI 316L stainless steel on biofilm adhesion in a seawater-cooled tubular heat exchanger-condenser. *Biofouling*, 32 (10), 1185–1193. doi: <https://doi.org/10.1080/08927014.2016.1241875>
- Maruschak, P., Panin, S., Chaurov, M., Bishchak, R., Polyvana, U. (2018). Effect of long-term operation on steels of main gas pipeline: Structural and mechanical degradation. *Journal of King Saud University - Engineering Sciences*, 30 (4), 363–367. doi: <https://doi.org/10.1016/j.jksues.2016.09.002>
- Whitehead, K. A., Verran, J. (2009). The Effect of Substratum Properties on the Survival of Attached Microorganisms on Inert Surfaces. *Marine and Industrial Biofouling*, 13–33. doi: https://doi.org/10.1007/978-3-540-69796-1_2
- An, Y. H., Dickinson, R. B., Doyle, R. J. (2000). Mechanisms of Bacterial Adhesion and Pathogenesis of Implant and Tissue Infections. *Handbook of Bacterial Adhesion*, 1–27. doi: <https://doi.org/10.1385/1-59259-224-4:1>
- Partington, E. (2006). *Stainless Steel in the Food & Beverage Industry. Materials and Applications*, 24.
- Jullien, C., Bénézech, T., Carpentier, B., Lebre, V., Faille, C. (2003). Identification of surface characteristics relevant to the hygienic status of stainless steel for the food industry. *Journal of Food Engineering*, 56 (1), 77–87. doi: [https://doi.org/10.1016/s0260-8774\(02\)00150-4](https://doi.org/10.1016/s0260-8774(02)00150-4)
- Dantas, L. C. de M., Silva-Neto, J. P. da, Dantas, T. S., Naves, L. Z., das Neves, F. D., da Mota, A. S. (2016). Bacterial Adhesion and Surface Roughness for Different Clinical Techniques for Acrylic Polymethyl Methacrylate. *International Journal of Dentistry*, 2016, 1–6. doi: <https://doi.org/10.1155/2016/8685796>
- Whitehead, K. A., Benson, P. S., Verran, J. (2015). Developing application and detection methods for *Listeria monocytogenes* and fish extract on open surfaces in order to optimize cleaning protocols. *Food and Bioproducts Processing*, 93, 224–233. doi: <https://doi.org/10.1016/j.fbp.2014.07.007>
- Whitehead, K. A., Verran, J. (2006). The Effect of Surface Topography on the Retention of Microorganisms. *Food and Bioproducts Processing*, 84 (4), 253–259. doi: <https://doi.org/10.1205/fbp06035>
- Hilbert, L. R., Bagge-Ravn, D., Kold, J., Gram, L. (2003). Influence of surface roughness of stainless steel on microbial adhesion and corrosion resistance. *International Biodeterioration & Biodegradation*, 52 (3), 175–185. doi: [https://doi.org/10.1016/s0964-8305\(03\)00104-5](https://doi.org/10.1016/s0964-8305(03)00104-5)
- Milledge, J. J. (2010). The cleanability of stainless steel used as a food contact surface: an updated short review. *Food Sci. Technol.*, 24 (3), 18–20.
- Bos, R., Mei, H. C., Gold, J., Busscher, H. J. (2000). Retention of bacteria on a substratum surface with micro-patterned hydrophobicity. *FEMS Microbiology Letters*, 189 (2), 311–315. doi: <https://doi.org/10.1111/j.1574-6968.2000.tb09249.x>
- Fadeeva, E., Truong, V. K., Stiesch, M., Chichkov, B. N., Crawford, R. J., Wang, J., Ivanova, E. P. (2011). Bacterial Retention on Superhydrophobic Titanium Surfaces Fabricated by Femtosecond Laser Ablation. *Langmuir*, 27 (6), 3012–3019. doi: <https://doi.org/10.1021/la104607g>
- Privett, B. J., Youn, J., Hong, S. A., Lee, J., Han, J., Shin, J. H., Schoenfisch, M. H. (2011). Antibacterial Fluorinated Silica Colloid Superhydrophobic Surfaces. *Langmuir*, 27 (15), 9597–9601. doi: <https://doi.org/10.1021/la201801e>
- Dou, X.-Q., Zhang, D., Feng, C., Jiang, L. (2015). Bioinspired Hierarchical Surface Structures with Tunable Wettability for Regulating Bacteria Adhesion. *ACS Nano*, 9 (11), 10664–10672. doi: <https://doi.org/10.1021/acsnano.5b04231>
- Cunha, A., Elie, A.-M., Plawinski L., Serro, A. P., Botelho do Rego, A. M., Almeida, A. et al. (2016). Femtosecond laser surface texturing of titanium as a method to reduce the adhesion of *Staphylococcus aureus* and biofilm formation. *Applied Surface Science*, 360, 485–493. doi: <https://doi.org/10.1016/j.apsusc.2015.10.102>
- Kukhtyn, M., Kravcheniuk, K., Beyko, L., Horiuk, Y., Skliar, O., Kernychnyi, S. (2019). Study of the influence of Savinase® Ecity 16L enzyme on biofilms formation of *staphylococcus aureus* on

- stainless steel with different roughness. EUREKA: Life Sciences, 2, 26–32. doi: <http://dx.doi.org/10.21303/2504-5695.2019.00858>
34. Vos, P., Garrity, G., Jones, D., Krieg, N. R. et. al. (Eds.) (2009). *Bergey's manual of systematic bacteriology. The Firmicutes*. Springer. doi: <https://doi.org/10.1007/978-0-387-68489-5>
35. Marchand, S., De Block, J., De Jonghe, V., Coorevits, A., Heyndrickx, M., Herman, L. (2012). Biofilm Formation in Milk Production and Processing Environments; Influence on Milk Quality and Safety. *Comprehensive Reviews in Food Science and Food Safety*, 11 (2), 133–147. doi: <https://doi.org/10.1111/j.1541-4337.2011.00183.x>
36. Kukhtyn, M., Vichko, O., Berhilevych, O., Horyuk, Y., Horyuk, V. (2016). Main microbiological and biological properties of microbial associations of «*Lactomyces tibeticus*». *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 7 (6), 1266–1272.

DOI: 10.15587/1729-4061.2019.162072

**STUDYING THE LOSS OF MASS BY CAULIFLOWER
DEPENDING ON AGROBIOLOGICAL FACTORS,
VARIETAL FEATURES, AND PACKAGE TECHNIQUE
(p. 22-31)**

Ludmila Pusik

Kharkiv Petro Vasylenko National Technical University
of Agriculture, Kharkiv, Ukraine

ORCID: <http://orcid.org/0000-0002-5465-2771>

Vladimir Pusik

Kharkiv Petro Vasylenko National Technical University
of Agriculture, Kharkiv, Ukraine

ORCID: <http://orcid.org/0000-0001-5028-9461>

Vitaliy Vlasovets

Kharkiv Petro Vasylenko National Technical University
of Agriculture, Kharkiv, Ukraine

ORCID: <http://orcid.org/0000-0002-6657-6761>

Ludmila Gaevaya

Kharkiv National Agrarian University named after V. V. Dokuchaev,
township Dokuchaevsky, Kharkiv region, Kharkiv district, Ukraine

ORCID: <http://orcid.org/0000-0002-8302-1776>

Artur Rozhkov

Kharkiv National Agrarian University named after V. V. Dokuchaev,
township Dokuchaevsky, Kharkiv region, Kharkiv district, Ukraine

ORCID: <http://orcid.org/0000-0001-9138-7973>

Nina Lyubymova

Kharkiv National Agrarian University named after V. V. Dokuchaev,
township Dokuchaevsky, Kharkiv region, Kharkiv district, Ukraine

ORCID: <http://orcid.org/0000-0001-8964-7326>

Iryna Safronska

Luhansk National Agrarian University, Kharkiv, Ukraine

ORCID: <http://orcid.org/0000-0003-1442-6312>

Oleksandr Romanashenko

Kharkiv Petro Vasylenko National Technical University
of Agriculture, Kharkiv, Ukraine

ORCID: <http://orcid.org/0000-0002-0857-6288>

Yevhen Kryshopt

Kharkiv National Agrarian University named after V. V. Dokuchaev,
township Dokuchaevsky, Kharkiv region, Kharkiv district, Ukraine

ORCID: <http://orcid.org/0000-0002-2868-3101>

Evgeniy Hryn

State Ecological Inspectorate in Kharkiv region, Kharkiv, Ukraine

ORCID: <http://orcid.org/0000-0002-4927-8228>

The aim of this study was to scientifically substantiate influence of weather conditions during vegetation season of cauliflower, the

content of some components of the chemical composition, physical properties, as well as packaging techniques, on the intensity of loss of cauliflower heads during storage. We established that the following types of loss occur during storage of cauliflower: weight loss due to a decrease in dry matter, weight loss due to partial evaporation of water, development of microorganisms and physiological disorders.

Weather conditions over the years of study were very variable. The sum of average daily temperatures ranged from 1,743.4 to 2,544.3 °C over the years of study, it exceeded the upper limit of the optimal amount by 58.9–380.3 °C. HTC (hydrothermal coefficient) was 0.58–1.10. Humidity was sufficient during the vegetation period. Conditions of the vegetation period affected weight loss of late-ripening cauliflower heads during storage in the following way: the greatest loss was 12.8–16.9 % in control in 2015, it was less in 2016 – 11.7–13.8 %, depending on a hybrid. The difference between hybrids over the years of study was significant. The natural decreases in weight loss of cauliflower heads during unpacked storage were for Casper F₁ hybrid – 13.7 %, for Santamaria F₁ – 12.7 %, for Skywalker F₁ – 15.5 % in 2015–2017.

The intensity of moisture reduction when cauliflower was stored at a temperature of 1±0.5 °C ranged from 0.27 to 9.3 %, depending on the packaging technique. The intensity of moisture evaporation in all hybrids was the same and it was 0.3 % per day under packaging by stretch film. Perforated stretch film increased the moisture evaporation slightly, from 0.37 % in Casper F₁ hybrid to 0.43 in Skywalker F₁ hybrid. The ratio of moisture loss to the loss of dry matter ranged from 11.2 in heads of Santamaria F₁ to 13.2 in heads of Casper F₁ during storage without packaging. And it was 0.23–0.29 when stored under SE, and 0.35–0.43 during storage under PSE.

We found out the correlation between cauliflower weight loss during storage on HTC, average daily temperature, and relative humidity of air in the vegetation period and on dry matter content, volume, and porosity of a head. The regression equations made it possible to forecast cauliflower weight loss during storage, depending on weather conditions of the vegetation period and on the dry matter content and physical indicators of heads.

Keywords: cauliflower, weight loss, weather conditions, water evaporation, types of packaging.

References

- Bondareva, L. L. (2013). New varieties and heterosis hybrids of cole crops of VNISSOK's breeding. *Vegetable crops of Russia*, 3, 32–33. doi: <https://doi.org/10.18619/2072-9146-2013-3-32-33>
- Houbein, L. (2017). *One Magic Square Vegetable Gardening: The Easy, Organic Way to Grow Your Own Food on a 3-Foot Square*. The Experiment, 352.
- Prusky, D. (2011). Reduction of the incidence of postharvest quality losses, and future prospects. *Food Security*, 3 (4), 463–474. doi: <https://doi.org/10.1007/s12571-011-0147-y>
- Koltunov, V. A. (2004). *Yakist plodoovochevoi produktsiyi ta tekhnolohiya yii zberihannia*. Ch. 1: Yakist i zberizhenist kartopli ta ovochiv. Kyiv: KNTEU, 568.
- Koltunov, V. A. (2002). *Prohnozuvannia yakosti prodovolchychk tovariv*. Kyiv, 199.
- Lohinov, M. I., Rosnovskiy, M. H., Lohinov, A. M. (2014). *Tekhnolohiya vyrobnytstva i pererobka produktiv roslinnytstva*. Hlukhiv: HNPU, 229.
- Cecilia, N., Emond, J.-P. (2007). Relationship between weight loss and visual quality of fruits and vegetables. *Proceedings of the Florida State Horticultural Society*, 120, 235–245.
- Paull, R. (1999). Effect of temperature and relative humidity on fresh commodity quality. *Postharvest Biology and Technology*, 15 (3), 263–277. doi: [https://doi.org/10.1016/s0925-5214\(98\)00090-8](https://doi.org/10.1016/s0925-5214(98)00090-8)
- Irtwange, S. V. (2006). Application of modified atmosphere packaging and related technology in postharvest handling of fresh fruits and veg-

- etables. *Agricultural Engineering International: CIGR Ejournal*, 8 (4). Available at: https://www.researchgate.net/profile/Simon_Irtwange/publication/228878406_Application_of_Modified_Atmosphere_Packaging_and_Related_Technology_in_Postharvest_Handling_of_Fresh_Fruits_and_Vegetables/links/54b642420cf28ebe92e7c0fd.pdf
10. Barabash, O. Yu., Taranenko, L. K., Sych, Z. D. Vplyv faktoriv navkolyshnoho seredovyscha na rist i rozvytok roslyn ovochevykh kultur. Available at: https://agromage.com/stat_id.php?id=216
 11. Pris, O. P., Kulik, A. S. (2015). Dynamika zminy masy zeleni petrushky pid chas zberihannia. *Khimiya kharchovykh produktiv i materialiv*, 2, 74–79. Available at: <http://oaji.net/articles/2016/1728-1456398475.pdf>
 12. Serdiuk, M. Ye., Stepanenko, D. S., Kiurchev, S. V. (2016). The study of mass loss intensity of plum fruits during storage. *Eastern-European Journal of Enterprise Technologies*, 1 (10 (79)), 42–48. doi: <https://doi.org/10.15587/1729-4061.2016.59694>
 13. Tan, W. L., Tao, L. R., Li, J. et. al. (201). Study on water loss of green cabbage under different relative humidity. *Science and Technology of Food Industry*, 7, 368–374.
 14. Kalashnik, V. F. (2011). Role of packaging in storing bell pepper. *Ovochivnytstvo i bashtannytstvo*, 57, 215–218.
 15. Barth, M. M., Zhuang, H. (1996). Packaging design affects antioxidant vitamin retention and quality of broccoli florets during postharvest storage. *Postharvest Biology and Technology*, 9 (2), 141–151. doi: [https://doi.org/10.1016/s0925-5214\(96\)00043-9](https://doi.org/10.1016/s0925-5214(96)00043-9)
 16. Kaynas, K., Özelkök, I. S. (1999). Effect of semperfresh on postharvest behavior of cucumber (*cucumis sativus* L.) And summer squash (*cucurbita pepo* L.) Fruits. *Acta Horticulturae*, 492, 213–220. doi: <https://doi.org/10.17660/actahortic.1999.492.27>
 17. Habibunnisa, Baskaran, R., Prasad, R. et. al. (2001). Storage behaviour of minimally processed pumpkin (*Cucurbita maxima*) under modified atmosphere packaging conditions. *European Food Research and Technology*, 212 (2), 165–169. doi: <https://doi.org/10.1007/s002170000211>
 18. Bezikonnyi, P. V. (2011). Vplyv pozakorenevoho vykorystannia mikroelementiv na zberezhennist koreneplodiv buriaka stolovoho. *Zbirnyk nakiu prats. Silskohospodarski nauky*, 19, 89–93. Available at: http://irbis-nbuv.gov.ua/cgi-bin/irbis_nbuv/cgiirbis_64.exe?C21COM=2&I21DBN=UJRN&P21DBN=UJRN&IMAGE_FILE_DOWNLOAD=1&Image_file_name=PDF/ZnpPdatu_2011_19_24.pdf
 19. Water loss and postharvest quality (2015). *Postharvest Technology of Horticultural Crops Short Course 2015*. 2015. Available at: <https://ucanr.edu/datastoreFiles/234-2858.pdf>
 20. Ghaouth, A., Arul, J., Ponnampalam, R., Boulet, M. (1991). Use of chitosan coating to reduce water loss and maintain quality of cucumber and bell pepper fruits. *Journal of Food Processing and Preservation*, 15 (5), 359–368. doi: <https://doi.org/10.1111/j.1745-4549.1991.tb00178.x>
 21. Sato, S., Kamiyama, M., Iwata, T., Makita, N., Furukawa, H., Ikeda, H. (2006). Moderate Increase of Mean Daily Temperature Adversely Affects Fruit Set of *Lycopersicon esculentum* by Disrupting Specific Physiological Processes in Male Reproductive Development. *Annals of Botany*, 97 (5), 713–738. doi: <https://doi.org/10.1093/aob/mcl037>
 22. Balan, E. F., Chumak, I. G., Kartofyanu, V. G., Iukuridze, E. Zh. (2007). Dinamika poter' plodovoovoshchnoy produktsii po etapam nepreryvnoy holodil'no-transportnoy cepi (HTC). *Holodil'shchik. Internet-vypusk*, 3 (27). Available at: http://www.holodilshchik.ru/index_holodilshchik_best_article_issue_3_2007.htm
 23. Zamorska, I. (2018). The effect duration of delay with cooling on natural mass losses of strawberry. *Pratsi Tavriyskoho derzhavnogo ahrotekhnolohichnoho universytetu*, 1 (18), 97–102.
 24. Lykhatskyi, V. I., Cherednychenko, V. M. (2010). Kapusta tsvitna. *Vynnytsia*, 167.
 25. Bublyk, M. O. (2002). Intehrlna otsinka pohodnykh faktoriv dlia vyroshchuvannia plodovykh kultur. *Visnyk aharnoi nauky*, 6, 31–33.
 26. Sharp, D. C., Townsend, M. S., Qian, Y., Shellhammer, T. H. (2014). Effect of Harvest Maturity on the Chemical Composition of Cascade and Willamette Hops. *Journal of the American Society of Brewing Chemists*, 72 (4), 231–238. doi: <https://doi.org/10.1094/asbcj-2014-1002-01>
 27. Parra-Coronado, A., Fischer, G., Camacho-Tamayo, J. H. (2015). Development and quality of pineapple guava fruit in two locations with different altitudes in Cundinamarca, Colombia. *Bragantia*, 74 (3), 359–366. doi: <https://doi.org/10.1590/1678-4499.0459>
 28. Díaz-Pérez, J. C., Muy-Rangel, M. D., Mascorro, A. G. (2006). Fruit size and stage of ripeness affect postharvest water loss in bell pepper fruit (*Capsicum annuum* L.). *Journal of the Science of Food and Agriculture*, 87 (1), 68–73. doi: <https://doi.org/10.1002/jsfa.2672>
 29. Becker, B. R., Fricke, B. A. (1996). Transpiration and respiration of fruits and vegetables. *Science et Technique du Froid*.
 30. Zamorskaya, I. L. (2017). Vliyanie razmerov yagod zemlyaniki na velichinu ubyli massy pri hranenii. *Pererabotka i upravlenie kachestvom sel'skohozyaystvennoy produktsii: sbornik statey III Mezhdunarodnoy nauchno-prakticheskoy konferencii*. Minsk: BGATU, 101–103.
 31. Bondarenko, H. L., Yakovenko, K. I. (Eds.) (2001). *Metodyka doslidnoi spravy v ovochivnytstvi i bashtannytstvi*. Kharkiv: Osnova, 369.
 32. Thompson, A. K. (2003). *Fruit and vegetables: harvesting, handling and storage*. Blackwell Publishing Ltd.
 33. Osokina, N. M., Kostetska, K. V. (2013). Physical and thermophysical properties of fruit green-stuffs depending on sort and degree of ripeness. *Zbirnyk naukovykh prats Umanskoho natsionalnoho universytetu sadivnytstva*, 83, 38–43.
 34. Getinet, H., Seyoum, T., Woldetsadik, K. (2008). The effect of cultivar, maturity stage and storage environment on quality of tomatoes. *Journal of Food Engineering*, 87 (4), 467–478. doi: <https://doi.org/10.1016/j.jfoodeng.2007.12.031>
 35. Koltunov, V. A., Puzik, L. M. (2004). Zberihannia harbuzovykh plodiv. Kharkiv, 365.
 36. Koltunov, V. A., Puzik, L. M. (2009). The structure of squash mass losses during storage. *Ovochivnytstvo i bashtannytstvo*, 55, 134–139.
 37. Koltunov, V. A., Pusik, L. M., Vakulenko, L. M. (2006). Vplyv rozmiru ploda na zberezhennist kabachkiv, dyni, ohiriv. *Sbornik nauchnykh rabot Krymskogo gosudarstvennogo agrarnogo universiteta*, 93, 56–60.
 38. Hu, W., Jiang, A., Qi, H., Pang, K., Fan, S. (2007). Effects of initial low oxygen and perforated film package on quality of fresh-cut cabbages. *Journal of the Science of Food and Agriculture*, 87 (11), 2019–2025. doi: <https://doi.org/10.1002/jsfa.2944>
 39. Bielinska, Ye. V. (2012). *Tryvale zberihannia koreneplodiv redysky: naukove obgruntuvannia, praktychne zastosuvannia*. Poltava: RVV PUET, 112.
 40. Osokina, N. M., Kostetska, K. V. (2013). Influence terms of storage is on the losses mass of fruit green-stuffs. *Visnyk aharnoi nauky Prychornomoria*, 1, 130–133.

DOI: 10.15587/1729-4061.2019.160220

SEPARATION OF TERPENES FROM LEMON ESSENTIAL OIL BY SELECTIVE FRACTIONATION UNDER A VACUUM (p. 32-36)

Natalya Frolova

National University of Food Technologies, Kyiv, Ukraine
ORCID: <http://orcid.org/0000-0002-5009-6026>

Anatoliy Ukrainets

National University of Food Technologies, Kyiv, Ukraine
ORCID: <http://orcid.org/0000-0001-7115-8006>

Iryna Sylka

National University of Food Technologies, Kyiv, Ukraine
ORCID: <http://orcid.org/0000-0002-2867-7414>

Aleksandra Nemirich

National University of Food Technologies, Kyiv, Ukraine

ORCID: <http://orcid.org/0000-0003-4731-3939>

Oleg Kuzmin

National University of Food Technologies, Kyiv, Ukraine

ORCID: <http://orcid.org/0000-0001-9321-6684>

The paper reports basic modern methods for isolating terpenes, outlines the advantages and reasons for this particular technological processing of essential oils, specifically lemon oil. The qualitative composition and quantitative ratios of components in lemon essential oil were determined by gas chromatography on a packed column (the stationary phase-dinonylphthalate). Based on the chromatographic profile, it was found that the key component of the oil is monoterpene d-limonene (70.60±4.61 %), which masks a harmonious manifestation of lemon flavor.

The paper reports results from theoretical calculations of selective fractionation under a vacuum that have been tested at a laboratory installation for the fractional distillation of essential oils, which simulates a full cycle of fractionation. The basic nodes of the installation were aligned with the schemes of universal plants that are common in industry.

The optimal modes for the selection of terpenes and terpenoid fractions are given, namely the value for the limit and working reflux number, pressure, and boiling temperature ranges.

The fraction of terpene, selected at a cube temperature from 67 °C to 70 °C, under a pressure of 2.64 kPa at a reflux ratio of 1:3, makes up 30.0 % by weight. The fraction of terpenoids, selected at a cube temperature from 84 °C to 96 °C, under a residual pressure from 0.33 to 0.66 kPa and at a reflux ratio of 1:14, makes up 60.0 % by weight.

Following the organoleptic study, it was established that the terpene fraction has a lemon flavor, due to the high content (80.0 %) of limonene, with notes of bergamot (β -myrcene – 5.12 %). Among other terpenoids, citral adds a muscat tone to the aroma, which, together with geraniol and linalool, acquires a noble aroma of citrus with a fine floral note.

All fractions of essential oils should be used in the food and cosmetic technology because the fraction of terpenes could be a valuable component in the creation of original composite fragrances.

Keywords: lemon essential oil, fractionation, deterpenation, gas chromatography, packed column, limonene.

References

1. Stuart, G. R., Lopes, D., Oliveira, J. V. (2001). Deterpenation of Brazilian orange peel oil by vacuum distillation. *Journal of the American Oil Chemists' Society*, 78 (10), 1041–1044. doi: <https://doi.org/10.1007/s11746-001-0385-x>
2. Arce, A., Marchiaro, A., Martínez-Ageitos, J. M., Soto, A. (2008). Citrus Essential Oil Deterpenation by Liquid-Liquid Extraction. *The Canadian Journal of Chemical Engineering*, 83 (2), 366–370. doi: <https://doi.org/10.1002/cjce.5450830226>
3. Gironi, F., Maschietti, M. (2008). Continuous countercurrent deterpenation of lemon essential oil by means of supercritical carbon dioxide: Experimental data and process modelling. *Chemical Engineering Science*, 63 (3), 651–661. doi: <https://doi.org/10.1016/j.ces.2007.10.008>
4. Arche, A., Soto, A. (2008). Citrus essential oils: Extraction and Deterpenation. *Tree And Forestry Science And Biotechnology*, 2 (1). Available at: <https://pdfs.semanticscholar.org/c10e/c1b0fe4229b7d407abaa150e05c6e2b2c168.pdf>
5. Fantin, G., Fogagnolo, M., Maietti, S., Rossetti, S. (2010). Selective Removal of Monoterpenes from Bergamot Oil by Inclusion in Deoxycholic Acid. *Journal of Agricultural and Food Chemistry*, 58 (9), 5438–5443. doi: <https://doi.org/10.1021/jf100072r>
6. Kuzmin, O., Suikov, S., Niemirich, O., Ditrich, I., Sylka, I. (2017). Effects of the water desalting by reverse osmosis on the process of formation of water-alcohol mixtures. 1H NMR spectroscopy studies. *Ukrainian Food Journal*, 6 (2), 239–257. doi: <https://doi.org/10.24263/2304-974x-2017-6-2-6>
7. Zheljazkov, V. D., Astatkie, T., Jeliaskova, E. A., Tatman, A. O., Schlegel, V. (2013). Distillation time alters essential oil yield, composition and antioxidant activity of female *Juniperus scopulorum* trees. *Journal of Essential Oil Research*, 25 (1), 62–69. doi: <https://doi.org/10.1080/10412905.2012.744704>
8. Gonçalves, D., Teschke, M. E. E., Koshima, C. C., Gonçalves, C. B., Oliveira, A. L., Rodrigues, C. E. C. (2015). Fractionation of orange essential oil using liquid-liquid extraction: Equilibrium data for model and real systems at 298.2K. *Fluid Phase Equilibria*, 399, 87–97. doi: <https://doi.org/10.1016/j.fluid.2015.04.022>
9. Cannon, J. B., Cantrell, C. L., Astatkie, T., Zheljazkov, V. D. (2013). Modification of yield and composition of essential oils by distillation time. *Industrial Crops and Products*, 41, 214–220. doi: <https://doi.org/10.1016/j.indcrop.2012.04.021>
10. Fang, T., Goto, M., Sasaki, M., Hirose, T. (2004). Combination of Supercritical CO₂ and Vacuum Distillation for the Fractionation of Bergamot Oil. *Journal of Agricultural and Food Chemistry*, 52 (16), 5162–5167. doi: <https://doi.org/10.1021/jf049895f>
11. Frolova, N., Silka, I. (2015). Technology of natural flavours from home raw material and apparatus base. *Novi rishennia v suchasnykh tekhnolohiyakh: Visnyk NTU «KhPI»*, 46. Available at: <http://dspace.nuft.edu.ua/jspui/handle/123456789/23482>
12. Carev, H. I., Carev, V. I., Katrakov, I. B. (2000). *Prakticheskaya gazovaya hromatografiya*. Barnaul: Izd-vo Alt. un-ta, 156.
13. Frolova, N., Ukrainets, A., Silka, I. (2009). Imitated distillation method a rational basis of essential oil fractionation. *Kharchova promyslovist*, 8, 96–99. Available at: <http://dspace.nuft.edu.ua/jspui/handle/123456789/478>
14. Morachevskiy, A. G., Smirnova, N. A., Piotrovskaya, E. M. et. al.; Morachevskogo, A. G. (Ed.) (1989). *Termodinamika ravnovesiya zhidkost'-par*. Leningrad: Himiya, 344.
15. Tadé, M. O., Bisowarno, B. H., Tian, Y.-C. (2004). Modelling and Control of Reactive Distillation Systems. *IFAC Proceedings Volumes*, 37 (1), 31–38. doi: [https://doi.org/10.1016/s1474-6670\(17\)38706-2](https://doi.org/10.1016/s1474-6670(17)38706-2)
16. Bagaturov, S. A. (1974). *Teoriya i raschet peregonki i rektifikatsii*. Moscow: Gostoptekhizdat, 435.
17. Silka, I. (2016). Evaluating the diet of Ukrainian military forces. *Molodyi vchenyi*, 5, 32–42. Available at: <http://dspace.nuft.edu.ua/jspui/handle/123456789/23536>
18. Frolova, N., Ukrainets, A., Sylka, I., Naumenko, K., Chepel, N. (2017). Actuality and ways of processing lokal spicy-aromatic raw materials into food flavors. *Scientific Works of National University of Food Technologies*, 23 (5 (2)), 220–228. doi: <https://doi.org/10.24263/2225-2924-2017-23-5-2-28>

DOI: 10.15587/1729-4061.2019.160186

ELECTROPHORETIC SYSTEM FOR EXPRESS

ANALYSIS OF WHEY PROTEIN FRACTIONS (p. 37-44)

Volodymyr Yukalo

Ternopil Ivan Puluj National Technical University, Ternopil, Ukraine

ORCID: <http://orcid.org/0000-0003-3553-6713>

Kateryna Datsyshyn

Ternopil Ivan Puluj National Technical University, Ternopil, Ukraine

ORCID: <http://orcid.org/0000-0002-9229-9059>

Liudmyla Storozh

Ternopil Ivan Puluj National Technical University, Ternopil, Ukraine

ORCID: <http://orcid.org/0000-0002-3635-2900>

Fractional specificity of biological action and ability to form bioactive peptides, which have a positive effect on different physiologi-

cal systems of the body in the processes of proteolysis and digestion, are characteristic for whey proteins. Prospects for the production and application of whey protein fractions are related to the necessity of their composition control.

The comparative analysis of the electrophoretic systems previously used for the milk protein analysis was carried out for the creation of an express analysis method of whey protein fractions. These are the anode disc electrophoresis system in the presence of sodium dodecyl sulfate, the Davis disc electrophoresis system for acidic proteins in native conditions, the system in a homogeneous polyacrylamide gel with urea. The Davis disc electrophoresis system for acidic proteins was chosen as the basis. For the adaptation of this system to the requirements of express analysis, the stacking polyacrylamide gel was removed from its composition and the concentration of the separating gel was reduced. The difference in the composition of electrode buffer and gel buffer ions was used to provide the high separation efficiency of protein fractions. This allows saving the effect of protein concentration in the whey sample on the first stages of electrophoresis. The position of the basic whey protein fractions on electrophoregrams was established with the help of homogeneous marker proteins (β -lactoglobulin and whey albumin).

An accessible electrophoresis system in the slabs of a homogeneous polyacrylamide gel for serial express analysis of the fractional composition of whey proteins has been proposed as a result of researches. The system allows reliable identification of four protein fractions (α -LA, β -LG, BSA and IG). Close average values and standard deviations of the content of these fractions in 15 whey samples of one milk batch, obtained by the densitometry of three electrophoregrams: β -LG (37.3 ± 4.2 , 36.5 ± 2.8 ; 38.3 ± 2.7), α -LA (15.8 ± 1.5 , 15.8 ± 1.3 , 16.4 ± 1.1), BSA (8.2 ± 1.1 , 8.0 ± 1.0 , 9.4 ± 1.1), IG (17.6 ± 1.9 , 17.4 ± 1.5 , 16.8 ± 1.5) testify about good reproducibility of the method. The proposed method may be useful for the express identification of the basic whey protein fractions, which are precursors of biologically active peptides.

Keywords: milk whey protein fractions, polyacrylamide gel electrophoresis, express analysis, densitometry.

References

- Hramcov, A. G. (2011). Fenomen molochnoy syvorotki. Sankt-Peterburg: Professiya, 804.
- Korhonen, H. (2009). Milk-derived bioactive peptides: From science to applications. *Journal of Functional Foods*, 1 (2), 177–187. doi: <https://doi.org/10.1016/j.jff.2009.01.007>
- Iukalo, A. V., Datsyshyn, K. Ye., Yukalo, V. G. (2013). Bioactive peptides of the cow milk whey proteins (*Bos Taurus*). *Biotechnologia Acta*, 6 (5), 49–61. doi: <https://doi.org/10.15407/biotech6.05.049>
- Le, T. T., Zhao, D., Larsen, L. B. (2019). Analytical Methods for Measuring or Detecting Whey Proteins. *Whey Proteins*, 155–184. doi: <https://doi.org/10.1016/b978-0-12-812124-5.00005-9>
- Farrell, H. M., Jimenez-Flores, R., Bleck, G. T., Brown, E. M., Butler, J. E., Creamer, L. K. et. al. (2004). Nomenclature of the Proteins of Cows' Milk – Sixth Revision. *Journal of Dairy Science*, 87 (6), 1641–1674. doi: [https://doi.org/10.3168/jds.s0022-0302\(04\)73319-6](https://doi.org/10.3168/jds.s0022-0302(04)73319-6)
- Neelima, Sharma, R., Rajput, Y. S., Mann, B. (2013). Chemical and functional properties of glycomacropeptide (GMP) and its role in the detection of cheese whey adulteration in milk: a review. *Dairy Science & Technology*, 93 (1), 21–43. doi: <https://doi.org/10.1007/s13594-012-0095-0>
- Le, T. T., Deeth, H. C., Larsen, L. B. (2017). Proteomics of major bovine milk proteins: Novel insights. *International Dairy Journal*, 67, 2–15. doi: <https://doi.org/10.1016/j.idairyj.2016.11.016>
- Marnila, P., Korhonen, H. (2011). Milk Proteins: Immunoglobulins. *Encyclopedia of Dairy Sciences*, 807–815. doi: <https://doi.org/10.1016/b978-0-12-374407-4.00436-2>
- Korhonen, H., Marnila, P. (2011). Milk Proteins: Lactoferrin. *Encyclopedia of Dairy Sciences*, 801–806. doi: <https://doi.org/10.1016/b978-0-12-374407-4.00435-0>
- Brandelli, A., Daroit, D. J., Corrêa, A. P. F. (2015). Whey as a source of peptides with remarkable biological activities. *Food Research International*, 73, 149–161. doi: <https://doi.org/10.1016/j.foodres.2015.01.016>
- Athira, S., Mann, B., Saini, P., Sharma, R., Kumar, R., Singh, A. K. (2014). Production and characterisation of whey protein hydrolysate having antioxidant activity from cheese whey. *Journal of the Science of Food and Agriculture*, 95 (14), 2908–2915. doi: <https://doi.org/10.1002/jsfa.7032>
- De Simone, C., Picariello, G., Mamone, G., Stiuso, P., Dicitore, A., Vanacore, D. et. al. (2009). Characterisation and cytomodulatory properties of peptides from Mozzarella di Bufala Campana cheese whey. *Journal of Peptide Science*, 15 (3), 251–258. doi: <https://doi.org/10.1002/psc.1093>
- Fox, P. F., Uniacke-Lowe, T., McSweeney, P. L. H., O'Mahony, J. A. (2015). *Dairy Chemistry and Biochemistry*. Springer, 584. doi: <https://doi.org/10.1007/978-3-319-14892-2>
- Basch, J. J., Douglas, F. W., Procino, L. G., Holsinger, V. H., Farrell, H. M. (1985). Quantitation of Caseins and Whey Proteins of Processed Milks and Whey Protein Concentrates, Application of Gel Electrophoresis, and Comparison with Harland-Ashworth Procedure. *Journal of Dairy Science*, 68 (1), 23–31. doi: [https://doi.org/10.3168/jds.s0022-0302\(85\)80792-x](https://doi.org/10.3168/jds.s0022-0302(85)80792-x)
- Yukalo, A., Yukalo, V., Shynkaryk, M. (2009). Electrophoretic separation of the milk protein. *Proceeding of the International Conference on Bio and Food Electrotechnologies*. Compiegne, 227–231.
- Iukalo, A. V. (2015). Identification of protein fractions of milk cows casein complex. *The Ukrainian Biochemical Journal*, 87 (4), 87–91. doi: <https://doi.org/10.15407/ubj87.04.087>
- Yukalo, V. G. (2005). Obtaining of casein protein complex fractions from cow milk. *Nutracos*, 5, 17–19.
- Yukalo, V., Datsyshyn, K. (2019). Gel filtration of cow milk whey proteins. *Food Science and Technology*, 12 (4), 72–78. doi: <https://doi.org/10.15673/fst.v12i4.1183>
- Yukalo, V. H., Yavorskyi, B. I., Storozh, L. A., Solovodzinska, I. Ye. (2007). Kilkisnyi elektroforetychnyi analiz bilkiv kazeinovoho kompleksu. *Biolojiya tvaryn*, 9 (1-2), 269–272.
- Skalka, V. V., Savchuk, O. M., Ostapchenko, L. I. (2010). Vyvchennia riznykh form kazeinu u molotsi metodom dysk-elektroforezu. *Fizyka zhyvoho*, 18 (3), 36–38.
- Osterman, L. A. (1981). *Metody issledovaniya belkov i nukleinovyyh kislot: Elektroforez i ul'tracentrifugirovanie*. Moscow: Nauka, 288.

DOI: 10.15587/1729-4061.2019.161855

FORMING THE STRUCTURE OF WHIPPED DESSERTS WHEN INTRODUCING THE FOOD ADDITIVE “MAGNETOFOOD” TO THEIR FORMULATION (p. 45-55)

Iryna Tsykhanovska

Ukrainian Engineering Pedagogics Academy, Kharkiv, Ukraine

ORCID: <http://orcid.org/0000-0002-9713-9257>

Viktoria Evlash

Kharkiv State University of Food Technology and Trade,

Kharkiv, Ukraine

ORCID: <http://orcid.org/0000-0001-7479-1288>

Alexandr Alexandrov

Ukrainian Engineering Pedagogics Academy, Kharkiv, Ukraine

ORCID: <http://orcid.org/0000-0003-3592-285X>

Barna Khamitova

M. Auezov South Kazakhstan State University, Shymkent, Kazakhstan
ORCID: <http://orcid.org/0000-0001-8377-3938>

Karyna Svidlo

Kharkiv Trade and Economics Institute of Kyiv National University
 of Trade and Economics, Kharkiv, Ukraine
ORCID: <http://orcid.org/0000-0002-0175-7756>

Olesia Nechuiviter

Ukrainian Engineering Pedagogics Academy, Kharkiv, Ukraine
ORCID: <http://orcid.org/0000-0003-2775-8471>

The food additive “Magnetofood”, in the form of a nanopowder with particles the size of (70–80) nm, has been designed and proposed as an improver for the structure of whipped desserts. “Magnetofood” can both independently form the structural and mechanical properties of whipped masses and influence a gel-forming agent, participating in chemical and electrostatic interactions with it. Therefore, the food additive “Magnetofood” can simultaneously influence several technological properties in the food disperse system: it can become a stabilizer, thickener, foam- and jelly-forming agents. Owing to Fe (II), nano dimensions, and the developed active surface, “Magnetofood” acquires sorption, complexing, emulsifying, moisture retaining, fat retaining, water-binding, stabilizing, structure-forming properties. That allows us to recommend “Magnetofood” as an additive with a comprehensive effect in order to form the structure of whipped desserts and to improve the quality of whipped dessert products. It has been established that introducing the additive “Magnetofood” to the samples of berry-fruit mousses and fruit-and-egg white jellies in the amount of 0.10 %, 0.15 %, 0.20 % by weight of the formulation mixture improves the structural-mechanical properties of whipped desserts. Using the additive “Magnetofood” decreases density by (29 ± 1) kg/m³ – for mousses, by (26 ± 1) kg/m³ – for fruit-and-egg white jellies, as well as the duration of whipping by ~15 % compared to control. It has been proven that introducing the additive “Magnetofood” contributes to an increase in: plastic strength – by 1.23 times; porosity – by (14.3 ± 0.7) % for mousses and by (12.7 ± 0.6) % – for fruit-and-egg white jellies; foam stability by (22.5 ± 1.1) % compared to control. In addition, a foam-forming capacity increases by (40 ± 2) % for mousses and by (55 ± 3) % for fruit-and-egg white jellies; effective viscosity – by (4.4 ± 0.2) % for mousses and by (4.1 ± 0.2) % – for fruit-and-egg white jellies compared to control. We have established that the rational content of the food additive “Magnetofood” equals 0.15 % of formulation composition.

The obtained experimental data could be applied when developing technologies for whipped dessert products.

Keywords: food additive “Magnetofood”, whipped desserts, stabilizing and structure-forming properties.

References

- Horichenko, N. V., Horalchuk, A. B., Omelchenko, S. B. (2014). Rol PAR u protsesakh formuvannya mizhfaznykh adsorbtsiynykh shariv pinoemulsiynykh produktiv. Innovatsiyni tekhnolohiyi rozvytku u sferi kharchovykh vyrobnytstv, hotelno-restorannoho biznesu, ekonomiky ta pidpriemnytstva: naukovi poshuky molodi: vseukr. nauk.-prakt. konf. molodykh uchenykh i studentiv. Ch. 1. Kharkiv: KhDUKhT, 5.
- Myronov, O. Yu., Horalchuk, A. B., Tovma, L. F. (2013). Obgruntuvannya vykorystannia poverkhnevo-aktyvnykh rehovyn v tekhnolohiyi pinopodibnoi produktsiyi na osnovi yaiechnoho bilka. Aktualni problemy rozvytku kharchovykh vyrobnytstv, restoran-noho ta hotelnogo hospodarstva i torhivli: vseukr. nauk.-prakt. konf. molodykh uchenykh i studentiv. Ch. 1. Kharkiv: KhDUKhT, 35.
- Omelchenko, S. B., Shania, I. M., Horalchuk, A. B. (2013). Obgruntuvannya vmistu emulhatoriv u skladi pinoemulsiynykh produktiv. Aktualni problemy rozvytku kharchovykh vyrobnytstv, restoran-noho hospodarstva i torhivli: vseukr. nauk.-prakt. konf. molodykh uchenykh i studentiv. Ch. 1. Kharkiv: KhDUKhT, 53.
- Omelchenko, S. B., Horalchuk, A. B. (2015). Vyznachennia roli poverkhnevo-aktyvnykh rehovyn u formuvanni mizhfaznykh adsorbtsiynykh shariv. Innovatsiyni aspekty rozvytku obladnannia kharchovoi i hotelnoi industrii v umovakh suchasnosti: mizhnar. nauk.-prakt. konf. Kharkiv: KhDUKhT, 291–292.
- Dickinson, E. (2006). Interfacial Particles in Food Emulsions and Foams. Colloidal Particles at Liquid Interfaces, 298–327. doi: <https://doi.org/10.1017/cbo9780511536670.009>
- Percevoy, F. V., Foshchan, A. L., Savgira, Yu. A. et. al. (2003). Proizvodstvo zheleynoy i vzbivnoy produktsii s ispol'zovaniem modifikatorov. Dnepropetrovsk: Porogi, 204.
- Ilyukha, N. G., Barsova, Z. V., Kovalenko, V. A., Tsykhanovska, I. V. (2010). Production technology and quality indices of a food additive based on magnetite. Eastern-European Journal of Enterprise Technologies, 6 (10 (48)), 32–35. Available at: <http://journals.urau.ua/eejet/article/view/5847/5271>
- Tsykhanovska, I., Evlash, V., Alexandrov, A., Lazarieva, T., Svidlo, K., Gontar, T. et. al. (2018). Substantiation of the mechanism of interaction between biopolymers of ryeandwheat flour and the nanoparticles of the magnetofood food additive in order to improve moistureretaining capacity of dough. Eastern-European Journal of Enterprise Technologies, 2 (11 (92)), 70–80. doi: <https://doi.org/10.15587/1729-4061.2018.126358>
- Tsykhanovska, I., Evlash, V., Alexandrov, A., Lazarieva, T., Bryzyska, O. (2018). Substantiation of the interaction mechanism between the lipo- and glucoproteids of rye-wheat flour and nanoparticles of the food additive «Magnetofood». Eastern-European Journal of Enterprise Technologies, 4 (11 (94)), 61–68. doi: <https://doi.org/10.15587/1729-4061.2018.140048>
- Karim, A. A., Bhat, R. (2008). Gelatin alternatives for the food industry: recent developments, challenges and prospects. Trends in Food Science & Technology, 19 (12), 644–656. doi: <https://doi.org/10.1016/j.tifs.2008.08.001>
- Campbell, G. (1999). Creation and characterisation of aerated food products. Trends in Food Science & Technology, 10 (9), 283–296. doi: [https://doi.org/10.1016/s0924-2244\(00\)00008-x](https://doi.org/10.1016/s0924-2244(00)00008-x)
- Lazidis, A., Hancocks, R. D., Spyropoulos, F., Kreuß, M., Berrocal, R., Norton, I. T. (2016). Whey protein fluid gels for the stabilisation of foams. Food Hydrocolloids, 53, 209–217. doi: <https://doi.org/10.1016/j.foodhyd.2015.02.022>
- Green, A. J., Littlejohn, K. A., Hooley, P., Cox, P. W. (2013). Formation and stability of food foams and aerated emulsions: Hydrophobins as novel functional ingredients. Current Opinion in Colloid & Interface Science, 18 (4), 292–301. doi: <https://doi.org/10.1016/j.cocis.2013.04.008>
- Dickinson, E. (2015). Structuring of colloidal particles at interfaces and the relationship to food emulsion and foam stability. Journal of Colloid and Interface Science, 449, 38–45. doi: <https://doi.org/10.1016/j.jcis.2014.09.080>
- Murray, B. S., Durga, K., Yusoff, A., Stoyanov, S. D. (2011). Stabilization of foams and emulsions by mixtures of surface active food-grade particles and proteins. Food Hydrocolloids, 25 (4), 627–638. doi: <https://doi.org/10.1016/j.foodhyd.2010.07.025>
- Phawaphuthanon, N., Yu, D., Ngamnikom, P., Shin, I.-S., Chung, D. (2019). Effect of fish gelatine-sodium alginate interactions on foam formation and stability. Food Hydrocolloids, 88, 119–126. doi: <https://doi.org/10.1016/j.foodhyd.2018.09.041>
- Dabestani, M., Yeganehzad, S. (2019). Effect of Persian gum and Xanthan gum on foaming properties and stability of pasteurized fresh egg white foam. Food Hydrocolloids, 87, 550–560. doi: <https://doi.org/10.1016/j.foodhyd.2018.08.030>
- Foshchan, A. L. (2010). Rehuliuuvannya reolohichnykh ta strukturno-mekhanichnykh vlastyvostei zheleinykh vyrobiv ta napivfabrykativ

- na osnovi kombinovanykh system drahle utvoriuvachiv. *Khlibopekarska i kondyterska promyslovist Ukrainy*, 2, 29–30.
19. Ignatova, T. A., Podkorytova, A. V. (2014). Ispol'zovanie gidrogeley karraginanov v tekhnologii zhelelynykh produktov. Aktual'nye problemy osvoiniya biologicheskikh resursov mirovogo okeana: materialy III Mezhdunar. nauch.-tekhn. konf. Vladivostok: Dal'rybvtuz, 58–63.
 20. Ellis, A. L., Mills, T. B., Norton, I. T., Norton-Welch, A. B. (2019). The hydrophobic modification of kappa carrageenan microgel particles for the stabilisation of foams. *Journal of Colloid and Interface Science*, 538, 165–173. doi: <https://doi.org/10.1016/j.jcis.2018.11.091>
 21. Bovšková, H., Míková, K. (2011). Factors influencing egg white foam quality. *Czech Journal of Food Sciences*, 29 (4), 322–327. doi: <https://doi.org/10.17221/435/2010-cjfs>
 22. Cotterill, O. J., Chang, C. C., Mcbee, L. E., Heymann, H. (1992). Metallic Cations Affect Functional Performance of Spray-Dried Heat-Treated Egg White. *Journal of Food Science*, 57 (6), 1321–1321. doi: <https://doi.org/10.1111/j.1365-2621.1992.tb06846.x>
 23. Corry, J. E. L., James, C., James, S. J., Hinton, M. (1995). Salmonella, Campylobacter and Escherichia coli O157:H7 decontamination techniques for the future. *International Journal of Food Microbiology*, 28 (2), 187–196. doi: [https://doi.org/10.1016/0168-1605\(95\)00056-9](https://doi.org/10.1016/0168-1605(95)00056-9)
 24. Kilasoniya, K. G. (2004). Using feijoa and kiwi puree for production of whipped confectionary products. *Pishchevaya promyshlennost'*, 12, 79.
 25. Iorgacheva, E. G. (2002). Pyure iz topinambura – recepturnyy ingredient konditerskikh izdeliy. *Zb. nauk. pr. ODAKhT*, 23, 120–124.
 26. Fioramonti, S. A., Perez, A. A., Aringoli, E. E., Rubiolo, A. C., Santiago, L. G. (2014). Design and characterization of soluble biopolymer complexes produced by electrostatic self-assembly of a whey protein isolate and sodium alginate. *Food Hydrocolloids*, 35, 129–136. doi: <https://doi.org/10.1016/j.foodhyd.2013.05.001>
 27. Mao, L., Boiteux, L., Roos, Y. H., Miao, S. (2014). Evaluation of volatile characteristics in whey protein isolate–pectin mixed layer emulsions under different environmental conditions. *Food Hydrocolloids*, 41, 79–85. doi: <https://doi.org/10.1016/j.foodhyd.2014.03.025>
 28. Wang, M.-P., Chen, X.-W., Guo, J., Yang, J., Wang, J.-M., Yang, X.-Q. (2019). Stabilization of foam and emulsion by subcritical water-treated soy protein: Effect of aggregation state. *Food Hydrocolloids*, 87, 619–628. doi: <https://doi.org/10.1016/j.foodhyd.2018.08.047>
 29. Kaprel'yanc, L. V., Iorgacheva, E. G., Banova, S. I. (2002). Modificirovannyye soeprodukty s uluchshennymi penoobrazuyushchimi i emul'giruyushchimi svoystvami. *Zernovi produkty i kombikormy*, 2, 23–25.
 30. Burgos-Díaz, C., Wandersleben, T., Olivos, M., Lichtin, N., Bustamante, M., Solans, C. (2019). Food-grade Pickering stabilizers obtained from a protein-rich lupin cultivar (AluProt-CGNA®): Chemical characterization and emulsifying properties. *Food Hydrocolloids*, 87, 847–857. doi: <https://doi.org/10.1016/j.foodhyd.2018.09.018>
 31. Jiang, X., Yucel Falco, C., Dalby, K. N., Siegmund, H., Arneborg, N., Risbo, J. (2019). Surface engineered bacteria as Pickering stabilizers for foams and emulsions. *Food Hydrocolloids*, 89, 224–233. doi: <https://doi.org/10.1016/j.foodhyd.2018.10.044>
 32. Hu, N., Wu, Z., Jin, L., Li, Z., Liu, W., Huang, D., Yang, C. (2019). Nanoparticle as a novel foam controller for enhanced protein separation from sweet potato starch wastewater. *Separation and Purification Technology*, 209, 392–400. doi: <https://doi.org/10.1016/j.seppur.2018.07.064>
 33. Zdobnov, A. I., Cyganenko, V. A. (2009). *Sbornik receptur blyud i kulinarnykh izdeliy: Dlya predpriyatiy obshchestvennogo pitaniya*. Kyiv: OOO "Izdatel'stvo Ariy", 680.
 34. Tsykhanovska, I., Yevlash, V., Alexandrov, A., Khamitova, B., Svidlo, K., Nechuiviter, O. (2019). Improving the technique of scrambled desserts using the food supplement "Magnetofood". *EUREKA: Life Sciences*, 2, 40–48. doi: <http://dx.doi.org/10.21303/2504-5695.2019.00856>
 35. Arkhipov, V. V., Ivanykova, T. V., Arkhipova, A. V. (2007). *Restoranna sprava: Asortyment, tekhnolohiya i upravlinnia yakistiu produktsiyi v suchasnomu restorani*. Kyiv: Firma «IIKOS», Tsentr navchalnoi literatury, 382.
 36. Zakharchuk, V. H., Kundilovska, T. A., Haidukovych, H. Ye. (2016). *Tekhnolohiya produktsiyi restorannoho hospodarstva*. Odessa: ONEU, Atlant VOI SOIU, 479.

DOI: 10.15587/1729-4061.2019.159983

ESTIMATION OF TECHNOLOGICAL PROPERTIES OF NUT MEALS AND THEIR EFFECT ON THE QUALITY OF EMULSION FOR BUTTER BISCUITS WITH LIQUID OILS (p. 56-64)

Elena Shydakova-Kameniuka

Kharkiv State University of Food Technology and Trade,
Kharkiv, UkraineORCID: <http://orcid.org/0000-0002-8550-7817>

Anna Novik

Oles Honchar Dnipro National University, Dnipro, Ukraine
ORCID: <http://orcid.org/0000-0003-4045-4878>

Yevhenii Zhukov

Kharkiv Trade and Economics Institute of Kyiv National University
of Trade and Economics, Kharkiv, Ukraine
ORCID: <http://orcid.org/0000-0002-8445-3490>

Yuliia Matsuk

Oles Honchar Dnipro National University, Dnipro, Ukraine
ORCID: <http://orcid.org/0000-0002-1899-3237>

Anna Zaparenko

Kharkiv Institute of Economics and Trade of Kyiv National University
of Economics and Trade, Kharkiv, Ukraine
ORCID: <http://orcid.org/0000-0001-9030-4128>

Philipp Babich

Dniprovskii University of the Humanities, Dnipro, Ukraine
ORCID: <http://orcid.org/0000-0003-1835-5021>

Svitlana Oliinyk

Kharkiv State University of Food Technology and Trade,
Kharkiv, Ukraine
ORCID: <http://orcid.org/0000-0003-4127-8247>

We have examined the technological properties of cedar nut meal (CNM) and walnut nut meal (WNM). It was established that nut meals are the highly dispersed powders whose degree of dispersion exceeds that of wheat flour. The size of 69 % of CNM is up to 40 μm, 72 % of WNM, and only 35 % of flour.

Compared with wheat flour, nut meals exposed to temperatures of 20...60 °C are characterized by better water-retaining capacity. When exposed to temperatures of 90 °C, meals and flour have close values for water-retaining capacity. It was noted that nut meals are better at emulsifying liquid fats than solid that are traditionally used in the technology of butter biscuits (margarine and butter). It is shown that meals are characterized by high fat-retention capacity (FRC) relative to liquid vegetable oil. It is noted that the value of FRC for nut meals in the temperature range 20...60 °C increases by 1.9 times; in the range 60...80 °C, FRC of the examined samples almost does not change; and when exposed to temperatures of 100...140 °C, it starts to decrease.

We have studied the quality of emulsions for butter biscuits for the case when 30 % of margarine are replaced with liquid vegetable oil while adding various amounts of nut meals. It was established that stability of the emulsion in which 30 % of margarine were replaced with liquid vegetable oil is 37.5 % less compared to control based on margarine. The introduction of CNM and WNM improves

the stability of such an emulsion. It was noted that the samples of emulsion with the addition of 40 % and 50 % of CNM and WNM in terms of the value for an indicator of stability are maximally close to the control whose fat base was margarine. This is confirmed by the results from studying the dispersion, the effective viscosity of emulsions, and by results from microscopy.

The obtained results are of practical significance in order to improve the technology of butter biscuits towards a partial substitution of margarine with liquid oils. That would improve the nutritional and biological value of butter biscuits.

Keywords: meal, walnut, cedar nut, water-retaining capacity, fat-retention capacity, fat-emulsifying capacity, emulsion, butter biscuits.

References

1. Za 2013-2017 gg prodazhi hlebobulochnykh i muchnykh konditerskikh izdeliy v mire vyrosli na 3,9%: so 103 do 107 mln t. Elektronnyi zhurnal «Biznes-analitik». Available at: <https://rusanalitic.ru/researches/za-gg-prodazhi-hlebobulochnykh-i-muchnykh-konditerskikh-izdeliy-v-mire-vyrosli-na-so-do-mln-t-1492>
2. Zanizdra, V. Vyrobnystvo pechyya. Baker-Group. Available at: <https://uk.baker-group.net/confectionery-formulations-technology-raw-materials-and-ingredients/4264-production-of-biscuits.html>
3. Carr, N. O., Hogg, W. F. (2005). A manufacturer's perspective on selected palm-based products. *Asia Pacific Journal of Clinical Nutrition*, 14 (4), 381–386.
4. Hotrum, N. E., Cohen Stuart, M. A., van Vliet, T., van Aken, G. A. (2004). Spreading of partially crystallized oil droplets on an air/water interface. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 240 (1-3), 83–92. doi: <https://doi.org/10.1016/j.colsurfa.2004.03.015>
5. Brun, M., Delample, M., Harte, E., Lecomte, S., Leal-Calderon, F. (2015). Stabilization of air bubbles in oil by surfactant crystals: A route to produce air-in-oil foams and air-in-oil-in-water emulsions. *Food Research International*, 67, 366–375. doi: <https://doi.org/10.1016/j.foodres.2014.11.044>
6. Bodnarchuk, O. V. (2015). Constructing of fatty basis of spread. *Zbirnyk naukovykh prats Vinnytskoho natsionalnoho ahrarnoho universytetu. Seriya: Tekhnichni nauky*, 2 (1 (89)), 31–36.
7. Zollner, N., Tato, F. (1992). Fatty acid composition of the diet: impact on serum lipids and atherosclerosis. *The Clinical Investigator*, 70 (11), 968. doi: <https://doi.org/10.1007/bf00180309>
8. Yildiz, F. (Ed.) (2010). *Advances in food biochemistry*. CRC Press Taylor & Francis Group. Available at: [http://gtu.ge/Agro-Lib/\[Fatih_Yildiz\]_Advances_in_Food_Biochemistry\(BookFi.org\).pdf](http://gtu.ge/Agro-Lib/[Fatih_Yildiz]_Advances_in_Food_Biochemistry(BookFi.org).pdf)
9. Cibizova, M. E., Myachina, A. G. (2005). Rasshirenie assortimenta zhirovyykh komponentov v recepturah muchnykh konditerskikh izdeliy. *Izvestiya vysshikh uchebnykh zavedeniy. Pishchevaya tekhnologiya*, 2-3, 67–69.
10. Tkachenko, A., Pakhomova, I. (2016). Consumer properties improvement of sugar cookies with fillings with non-traditional raw materials with high biological value. *Eastern-European Journal of Enterprise Technologies*, 3 (11 (81)), 54–61. doi: <https://doi.org/10.15587/1729-4061.2016.70950>
11. Bila, O., Kirpichenkova, O. (2017). Technology shortbread increased food value. *Teoriya i praktika srovennoy nauki*, 6 (24). Available at: <https://docplayer.ru/60913571-Tekhnologiya-pesochnogo-pechenya-povyshennoy-pishchevoy-cennosti.html>
12. Renzyaeva, T. V., Tuboltseva, A. S., Artyushina, S. I. (2015). Development of formulation and technology of gluten-free cookies based on natural vegetable raw materials. *Food Processing: Techniques and Technology*, 39 (4), 87–92.
13. Merman, A. D. (2013). Razrabotka i ochenka kachestva muchnykh konditerskikh izdeliy s rastitel'nyimi maslami. *Kemerovo*, 169.
14. Citrusovye volokna Herbacel AQ Plus – tip N. Specifikatsii dlya pishchevykh dobavok i receptury. Available at: <http://specin.ru/kletchatka/109.htm>
15. Renzyaeva, T. V., Merman, A. D. (2013). Modeling of the formula for butter biscuit with functional properties. *Tekhnika i tekhnologiya pishchevykh proizvodstv*, 1, 35–41.
16. Hordienko, L. V., Zhydetska, I. V. (2010). Vplyv spivvidnoshennia retsepturnykh komponentiv na reolohichni vlastyivosti emulsiyi dlia pishchoho tista. *Naukovi pratsi Odeskoi natsionalnoi akademiyi kharchovykh tekhnolohiyi*, 1 (38), 214–217.
17. Maner, S., Sharma, A. K., Banerjee, K. (2015). Wheat Flour Replacement by Wine Grape Pomace Powder Positively Affects Physical, Functional and Sensory Properties of Cookies. *Proceedings of the National Academy of Sciences, India Section B: Biological Sciences*, 87 (1), 109–113. doi: <https://doi.org/10.1007/s40011-015-0570-5>
18. Gorodyska, O., Grevtseva, N., Samokhvalova, O., Gubsky, S., Gavrish, T., Denisenko, S., Grigorenko, A. (2018). Influence of grape seeds powder on preservation of fats in confectionary glaze. *Eastern-European Journal of Enterprise Technologies*, 6 (11 (96)), 36–43. doi: <https://doi.org/10.15587/1729-4061.2018.147760>
19. Mihalik, V. S., Tkachenko, L. V. (2014). Innovation technology of cookies. *SWorld. Modern problems and ways of their solution in science, transport, production and education*. Available at: <https://sworld.com.ua/konfer35/612.pdf>
20. Yurchenko, S. L., Kolesnikova, M. B., Yurchenko, E. P. (2016). Improvement of prescription composition shortbread with using cellulose. *Molodyi vchenyi*, 3 (30), 284–287.
21. Korkach, G. V., Krusir, G. V., Egorova, A. V., Kushnir, G. (2015). Change in the quality of sugar cookies with the introduction of prebiotic supplement. *Kharchova nauka ta tekhnolohiya*, 9 (3), 49–56. doi: <https://doi.org/10.15673/2073-8684.3/2015.50282>
22. Samokhvalova, O. V., Chernikova, Yu. O., Oliinyk, S. H., Kasabova, K. R. (2015). The effect of microbial polysaccharides on the properties of wheat flour. *Eastern-European Journal of Enterprise Technologies*, 6 (10 (78)), 11–15. doi: <https://doi.org/10.15587/1729-4061.2015.56177>
23. Shidakova-Kamenyuka, E., Novik, A., Kasabova, K., Kravchenko, O. (2015). The perspectives of using extraction cake of nuts for the enrichment of floury confectionary products. *Prohresyvnii tekhnika ta tekhnolohiya kharchovykh vyrobnytstv, restorannoho hospodarstva ta torhivli*, 2, 69–81.
24. Shidakova-Kamenyuka, E., Novik, A., Bolhovitina, E. (2017). Analiz soderzhaniya osnovnykh pishchevykh veshchestv v produktakh pererabotki greckogo i kedrovogo orekha. *Scientific Letters of Academic of Michal Baludansky*, 5 (4), 121–124.
25. Subbotina, M. A. (2009). Technology of liquid milk plant mixtures for the ice cream. *Tekhnika i tekhnologiya pishchevykh proizvodstv*, 4, 31–35.
26. Zolotim, A. Yu., Simonenko, S. V. (2012). Mayonezy i mayoneznye sousy na osnove emul'sii yadra kedrovogo orekha. *Maslozhirovaya promyshlennost'*, 3, 15–17.
27. Orlova, O. Yu., Nasonova, Yu. K. (2014). Ispol'zovanie greckogo orekha molochno-voskovoy spelosti dlya razrabotki funktsional'nykh produktov pitaniya. *Nauchnyi zhurnal NIU ITMO. Seriya «Processy i apparaty pishchevykh proizvodstv»*, 2.
28. Hvylya, S. I. (2011). Mikrostruktura i sostav polufabrikatov dlya gerodieticheskogo pitaniya s myasom kur i zhmyhom kedrovogo orekha. *Myasnoy biznes*, 10, 63–65.
29. Lapshinskaya, N. A., Dragun, N. A., Alenicheva, Yu. Yu. (2009). Molochnye produkty s kombinirovannoy zhirovoy fazoy. *Sibirskiy torgovo-ekonomicheskii zhurnal*, 8, 163–165.
30. Rahbari, M., Aalami, M., Kashaninejad, M., Maghsoudlou, Y., Aghdaei, S. S. A. (2014). A mixture design approach to optimizing low cholesterol mayonnaise formulation prepared with wheat germ protein isolate. *Journal of Food Science and Technology*, 52 (6), 3383–3393. doi: <https://doi.org/10.1007/s13197-014-1389-4>

31. Cherevko, A., Kiptelaya, L., Mikhaylov, V., Zagorulko, A., Zagorulko, A. (2015). Development of energy-efficient IR dryer for plant raw materials. *Eastern-European Journal of Enterprise Technologies*, 4 (8 (76)), 36–41. doi: <https://doi.org/10.15587/1729-4061.2015.47777>
32. Shakerardekani, A., Karim, R., Ghazali, H., Chin, N. (2013). Textural, Rheological and Sensory Properties and Oxidative Stability of Nut Spreads – A Review. *International Journal of Molecular Sciences*, 14 (2), 4223–4241. doi: <https://doi.org/10.3390/ijms14024223>
33. Dudkin, M. S., Chernov, N. K., Kazanceva, I. S. et. al. (1988). *Pishchevye volokna*. Kyiv, 152.
34. Arhipov, A. N. (2009). Teoreticheskie osnovy strukturoobrazovaniya dispersnykh sistem dlya pridaniya im agregatnoy ustoychivosti. *Tekhnika i tekhnologiya pishchevykh proizvodstv*, 3, 17–19.
35. Sylchuk, T. A., Nazar, M. I., Karpenko, T. S., Dotsenko, V. F., Tsyrlukova, V. V. (2015). Structural and mechanical properties of rye wheat dough with peas cellulose. *Journal of Food Science and Technology*, 2, 86–89. doi: <https://doi.org/10.15673/2073-8684.31/2015.44280>
36. Kryvoruchko, M., Forostyana, N. (2016). Rheological properties of flour dough with coconut fiber. *Tovary i rynky*, 2, 177–184.
37. Awad-Allah, M. A. A. (2013). Evaluation of Selected Nuts and Their Proteins Functional Properties. *Journal of Applied Sciences Research*, 9 (1), 885–896.

DOI: 10.15587/1729-4061.2019.162671

STUDYING THE EFFECT OF THE INTEGRATED BREAD BAKING IMPROVER «MINERAL FRESHNESS SUPER» ON CONSUMER PROPERTIES OF WHEAT BREAD (p. 65-72)

Olena Bilyk

National University of Food Technologies, Kyiv, Ukraine
ORCID: <http://orcid.org/0000-0003-3606-1254>

Yulia Bondarenko

National University of Food Technologies, Kyiv, Ukraine
ORCID: <http://orcid.org/0000-0002-3781-5604>

Oksana Kochubei-Lytvynenko

National University of Food Technologies, Kyiv, Ukraine
ORCID: <http://orcid.org/0000-0003-0712-448X>

Esma Khalikova

National University of Food Technologies, Kyiv, Ukraine
ORCID: <http://orcid.org/0000-0001-5785-1306>

Albina Fain

Podilsky Special Educational and Rehabilitation Socio-Economic College, Kamianets-Podilsky, Ukraine
ORCID: <http://orcid.org/0000-0002-9107-7198>

Scientists from the National University of Food Technologies (Ukraine) developed the integrated bread baking improver “Mineral Freshness Super” whose formulation includes nutritional supplements with the GRAS status. The integrated bread baking improver is composed of a functional base, white pharmacopoeial clay, and an active part – the enzymatic preparations Alphamalt VC 5000 and Novamil 1500MG, maltodextrin, sunflower-derived fat-free lecithin, apple pectin, dry wheat gluten and ascorbic acid. Technological efficiency has been proven to use the integrated bread baking improver “Mineral Freshness Super” in the amount of 1.5 % by weight of flour in order to slow the staling of wheat bread, baked according to the accelerated technology.

We have determined regularities in the influence of the integrated bread baking improver “Mineral Freshness Super” on quality of wheat bread. It was established that introducing it to dough results in an increase in the relative volume of products, improves shape formation, porosity, and reduces the duration of fermentation by three times, specifically to 20 min.

It has been proven that products with the addition of the integrated bread baking improver “Mineral Freshness Super” retain freshness better, which is confirmed by the increase in the overall deformation of crumb, by a smaller layer under the crust, and by fewer layers of air in the products’ crumb. We have observed more accumulation of dextrins and bisulfite binding substances in products when using integrated bread baking improver “Mineral Freshness Super”, indicating the inhibition of staling and improvement in consumer properties.

Results of our study prove the expediency of using the integrated bread baking improver “Mineral Freshness Super” in the technology of wheat bread to prolong its freshness up to 72 h when storing unpacked.

Keywords: integrated bread baking improver, wheat bread, staling, under-the-crust layer, bound moisture.

References

1. Vasylenko, O. M. (2017). Stan khlibopekarskoi haluzi Ukrainy v suchasnykh umovakh. *Materialy mizhnarodnykh nauko-vo-praktychnykh konferentsiy «Innovatsiini tekhnolohiyi u vyrobnytsvi» ta «Zdobutky ta perspektyvy rozvytku kondyterskoi haluzi»*. Kyiv: NUKhT, 7–11.
2. Pashchenko, L. P., Zharkova, I. M. (2014). *Tekhnologiya hlebopekarnogo proizvodstva*. Sankt-Peterburg: Lan', 372.
3. Petuhov, M. M., Kolyada, E. V. (2013). Sovremennyye predstavleniya o svezhesti hlebobulochnykh izdeliy. *Sovremennaya trgovlya: teoriya, praktika, perspektivy razvitiya: Materialy Tret'ey mezhdunarodnoy innovatsionnoy nauchno-prakticheskoy konferentsii*. Moscow: Izdatel'stvo Moskovskogo gumanitarnogo universiteta, 308–311.
4. Nechaev, A. P., Kraus, S. V., Fihntner, E. et. al. (2013). Pishchevye ingredienty v proizvodstve hlebobulochnykh i muchnykh konditerskiykh izdeliy. Moscow: DeLi plyus, 527.
5. Koryachkina, S. Ya., Matveeva, T. V. (2013). Funktsional'nye pishchevye ingredienty i dobavki dlya hlebobulochnykh i konditers'kiykh izdeliy. Sankt-Peterburg: GIOR, 628.
6. Shevre, S. (2008). Kak sdelat' produkciyu luchshe. *Hlebopekarskoe i konditerskoe delo*, 2, 40–42.
7. Bilyk, O., Bondarenko, Y., Desyk, M., Sokolenko, A., Kovbasa, V., Bondar, V. (2018). Research into effectiveness of using the integrated bread baking improver «Mineral Freshness +» to slow down the staling of bakery products. *Eastern-European Journal of Enterprise Technologies*, 4 (11 (94)), 69–78. doi: <https://doi.org/10.15587/1729-4061.2018.140333>
8. Bilyk, O. A. (2017). Rozrobka kompleksnykh khlibopekarskykh polipshuvachiv dlia khlibobulochnykh vyrobiv podovzhenoho terminu zberihannia. *Naukovi pratsi NUKhT*, 23 (5), 239–247.
9. Matveeva, I. V. et. al. (2009). Kompleksnye pishchevye dobavki – hlebopekarnye uluchshiteli na osnove fermentov i vitaminov. *Hlebopechenie Rossii*, 5, 18–19.
10. Kitissu, P., Andreev, A. (2009). Ispol'zovanie fermentov v tekhnologii bystrozamarozhennykh testovykh polufabrikatov. *Hleboprodukty*, 4, 52–53.
11. Razumnyye innovatsii: novinki v assortimente kompanii Backaldrin (2013). *Hleboprodukty*, 1, 34–35.
12. Gamma Soft. Available at: https://zeelandia.ua/fileadmin/user_upload/goods/media/products/Gamma_Soft.pdf
13. Prodlevaem svezhest' hleba s uluchshitelem «Mazhimiks» (2009). *Mir produktov*, 4, 20–21.
14. Kraus, S. V. (2008). Predstavlyаем ООО «IREKS». Pishchevye ingredienty: syr'e i dobavki, 1, 6–7.
15. Danilyuk, E. (2011). Nuzhen vkus i aromat? Pomozhet «Puratos». *Hleboprodukty*, 9, 40–41.
16. Korshenko, L. (2014). Stabilization of wheat bread's quality with low baking properties. On-line Journal «Naukovedenie», 6. doi: <https://doi.org/10.15862/115tvn614>

17. Zyuz'ko, A. S., Korostova, E. V., Bondarenko, V. I. (2011). Razrabotka kompleksnogo uluchshatelya dlya povysheniya kachestva hleba iz pshenichnoy muki. *Izvestiya vuzov. Pishchevaya tekhnologiya*, 4, 24–25.
18. Korshenko, L. O., Chizhikova, O. G., Tanashkina, T. V., Docenko, S. M., Abdulaeva, N. N., Semenyuta, A. A. (2014). Obosnovanie ispol'zovaniya grechnevogo soloda pri razrabotke kompozicii hlebopekarnogo uluchshatelya. *Tekhnika i tekhnologiya pishchevykh proizvodstv*, 1, 49–53.
19. Lazovenko, I. A., Stepycheva, N. V. (2015). Use of enzymatic agent as constituent of complex bakery improver with anti-staling properties. *Trudy BGTU. Himiya, tekhnologiya organicheskikh veshchestv i biotekhnologiya*, 4, 198–201.
20. Acoski, T., Stanojeska, M. (2011). Influence of the enzyme complex in additive composition on bread yield in breadmaking process. *Journal of Hygienic Engineering and Design*, 243–246.
21. Mansurov, V., G. Galitsyn, Y., V. Malin, T., A. Teys, S., S. Zhuravlev, K., Cora, I., Pecz, B. (2018). Van der Waals and Graphene-Like Layers of Silicon Nitride and Aluminum Nitride. *2D Materials* [Working Title]. doi: <https://doi.org/10.5772/intechopen.81775>
22. Lebedenko, T. Ye., Pshenyshniuk, H. F., Sokolova, N. Yu. (2014). *Tekhnolohiya khlibopekarskoho vyrobnytstva. Praktykum*. Odessa: «Osvita Ukrainy», 392.
23. Stele, R. (2008). *Srok godnosti pishchevykh produktov. Raschet i ispytanie*. Sankt-Peterburg: Professiya, 480.
24. Drobot, V. I. (Ed.) (2015). *Tekhnokhimichniy kontrol syrovyny ta khlibobulochnykh i makaronnykh vyrobiv*. Kyiv: NUKhT, 902.
25. Kragh, K. (2003). Amylases in baking. *Recent Advances in Enzymes in Grain Processing*. Leuven, 221–227.
26. Nilova, L. P., Kalinina, I. V., Naumenko, N. V. (2013). Differential thermal analysis method to assess the quality of food products. *Vestnik YUUrGU. Seriya: Pishchevye i biotekhnologii*, 1, 43–48.
27. Popper, L. (2010). Enzymes: the best friends of flours. *Food Science*, 34–46.
28. Correa, M. J., Pérez, G. T., Ferrero, C. (2012). Pectins as Breadmaking Additives: Effect on Dough Rheology and Bread Quality. *Food and Bioprocess Technology*, 5 (7), 2889–2898. doi: <https://doi.org/10.1007/s11947-011-0631-6>
29. BeMiller, J. N. (2009). One Hundred Years of Commercial Food Carbohydrates in the United States. *Journal of Agricultural and Food Chemistry*, 57 (18), 8125–8129. doi: <https://doi.org/10.1021/jf8039236>
30. BeMiller, J., Whistler, R. (Eds.) (2009). *Starch: Chemistry and Technology*. Academic press: Burlington, 894.
31. Stauffer, C. E. (2000). Emulsifiers as antistaling agents. *Cereal Foods World*, 45 (3), 106–110.
32. Forsell, P., Shamekh, S., Härkönen, H., Poutanen, K. (1998). Effects of native and enzymatically hydrolysed soya and oat lecithins in starch phase transitions and bread baking. *Journal of the Science of Food and Agriculture*, 76 (1), 31–38. doi: [https://doi.org/10.1002/\(sici\)1097-0010\(199801\)76:1<31::aid-jsfa915>3.0.co;2-2](https://doi.org/10.1002/(sici)1097-0010(199801)76:1<31::aid-jsfa915>3.0.co;2-2)