

DESIGN OF ANTIGEN SYNTHESIS AND PREPARATION AND CHARACTERIZATION OF SPECIFIC AND EURYTROPIC ANTIBODIES AGAINST B-GROUP AFLATOXINS

Yanan Wang

postgraduate student

Sumy National Agrarian University (Sumy, Ukraine)

College of Animal Science and Veterinary Medicine, Henan Institute of Science and Technology (Xinxiang, China)

ORCID: 0000-0001-9537-2947

wyn564@126.com

Hanna Fotina

Doctor of Veterinary Sciences, Professor

Sumy National Agrarian University (Sumy, Ukraine)

ORCID: 0000-0002-0761-3681

hanna.fotina@snau.edu.ua

The aim of this study was to prepare B-group aflatoxins (BGAFs) antibody with strong specificity and good eurytopicity. According to the molecular structure and active site of aflatoxin B₁ (AFB₁), the BGAFs artificial antigen AFB₁-BSA was prepared by 6 methods such as oxime active ester (OAE), methylation of ammonia (MOA), mixed anhydride (MA), semi acetal (SA), epoxide (EP) and enol ether derivative (EED) and identified by UV and SDS-PAGE. Polyclonal antibodies against AFB₁ (AFB₁ pAb) were prepared by immunizing New Zealand rabbits with AFB₁-BSA, and the titers of AFB₁ pAb was detected by indirect ELISA, the sensitivity of AFB₁ pAb was analyzed by indirect competitive ELISA (icELISA) and the specificity and eurytopicity of AFB₁ pAb was analyzed by cross-reactivity (CR) test. The results showed that AFB₁-BSA was synthesized successfully and the best one was OAE method among 6 synthesis methods of BGAFs artificial antigen and its conjugation ratio of AFB₁ to BSA was about 8.46:1. The immune efficacy of OAE method was the best, its AFB₁ pAb had high titers of 1:1(28×10⁴) by indirect ELISA, a good sensitivity with the 50% inhibition concentration (IC₅₀) of 10.32 µg/L to AFB₁ by icELISA and a high CR to AFB₂ of 75.21%, AFG₁ of 44.13%, AFG₂ of 14.72%, AFM₁ of 16.36% and AFM₂ of 1.44%, respectively. In this study, AFB₁ pAbs with high titer, sensitivity, specificity and eurytopicity were prepared, which laid a matter and technical foundation for the establishment of BGAFs immunoassay.

Key word B-group aflatoxins, antigen synthesis design, polyclonal antibody, characteristics analysis

DOI: <https://doi.org/10.32845/bsnau.vet.2020.4.8>

Introduction. Aflatoxins (AFs) are a group of toxic secondary metabolites containing similar molecular structures (difuran ring and oxyheteronaphthalidone). They are produced by *Aspergillus flavus* and *Aspergillus parasiticus* through the polyketone pathway. At present, more than 20 members of the aflatoxin (AF) family have been found in food pollution. Among them, AF of group B (B group aflatoxins, BGAFs) have strong toxicity, wide pollution, and high content. Various toxic effects such as carcinogenicity, teratogenicity and immunosuppressiveness have become the main targets of food AF contamination detection (Sun D.D et al., 2015). BGAFs include AFB₁ and AFB₂, both of which are closely related to food pollution, and both exist at the same time, mainly AFB₁, AFB₂ pollution is accompanied by AFB₁, and has a toxic additive effect (Luo, X. et al., 2018). Therefore, there are two regulations for the maximum residue limits (MRLs) of BGAFs in foods. the current AFB₁ MRL standard of food and agricultural products in China is "GB 2761-2017 limit of fungal toxins in food" which One is that some countries including my country adopt AFB₁ MRLs, such as the current AFB₁ MRL standard of food and agricultural products in China is "GB 2761-2017 limit of fungal toxins in food" (CHINA. National Food Safety Standard Limit of mycotoxins in food. 2017), corn and its products ≤20 µg·kg⁻¹, rice and its products ≤10 µg·kg⁻¹, wheat and its products ≤5 µg·kg⁻¹. Second, some countries use MRLs for the total amount of BGAFs (B₁+B₂), such as EU ≤4 µg·kg⁻¹, Japan ≤10 µg·kg⁻¹, and US FDA ≤15 µg·kg⁻¹. There are many current analytical methods for food BGAFs contamination, mainly using instrumental analysis and

immunoassay. In particular, immunoassay has become a technology because of its strong specificity, high sensitivity, simple operation, large-scale screening and on-site detection. Indispensable technical means, the key to establishing a BGAFs immunoassay method is to obtain excellent antibodies, and hapten design and antigen synthesis are the prerequisites for preparing excellent antibodies (Gefen T. et. al., 2015). There have been related reports on the research of BGAFs antigen synthesis methods at home and abroad (Mongkon, W. et. al., 2017, Xiao L.W, et. al., 2017), but there are no reports on the design of different hapten molecules, antigen synthesis and comparative analysis of antibody characteristics. In this study, AFB₁ was used as the starting material for the reaction. Polyclonal antibodies (pAbs) were prepared through different AFB₁ hapten molecular design and antigen synthesis methods, and their characteristics were analyzed to screen out the best hapten and antigen synthesis methods. It lays the foundation for the preparation of high-quality monoclonal antibodies of BGAFs with high sensitivity, broad recognition spectrum and strong specificity (Zhou, Y. et. al., 2007).

Aim The aim of this study was to prepare B-group aflatoxins (BGAFs) antibody with strong specificity and good eurytopicity.

Materials and Methods

Main reagents, solutions and experimental animals

AFB₁, AFB₂, AFG₁, AFG₂ standard products, Singapore Pribolab product; Cationized bovine serum albumin (cBSA), goat anti-rabbit enzyme-labeled secondary antibody (GaRlgG-

HRP), American Sigma product. The diluent used in the enzyme-linked immunosorbent assay (ELISA) is 0.01 mol·L⁻¹ pH7.4 phosphate buffer solution (PBS); the washing solution is PBS containing 0.5 g·L⁻¹ Tween-20 (PBST); the blocking solution is PBST containing 50 g·L⁻¹ porcine serum; the coating solution is 0.1 mol·L⁻¹ carbonate buffer solution (CBS) with pH 9.6. The experimental animals were 18 male New Zealand white rabbits

at the age of 2 months and weighing 1±0.2 kg. They were provided by the Experimental Animal Center of Xinxiang Medical College. They were divided into 6 groups, each with 3 rabbits.

BGAFs artificial antigen synthesis design

According to the active sites on the molecular structure of AFB₁ (Figure 1), the following six methods are proposed to prepare artificial antigen AFB₁-BSA (Table 1).




Fig.1 Molecular structure of AFB

Table1

The hapten design and antigens synthesis of AFB₁

Antigen synthesis design					Antigen synthesis route	Reaction principle
Active site	Active group	Reaction method	Introduction group	Synthesis Method		
1	Carbonyl	Oximation	carboxyl	OAE		The oximation of the 1-position carbonyl group of the active site of AFB ₁ to AFB ₁ O, the introduction of the carboxyl active group, the active ester method under the action of the coupling agent dicyclohexylcarbodiimide (DCC), the synthesis of AFB ₁ O and BSA in the form of a single amide bond AFB ₁ -BSA [7,8].
2	Active hydrogen	Mannich	Aminomethyl	MOA		Using the 2-position α-active hydrogen of AFB ₁ , through Mannich reaction, the α-active hydrogen and the amino group of BSA undergo an aminomethylation reaction, which is coupled in the form of Mannich base to synthesize AFB ₁ -BSA [9,10].

Antigen synthesis design					Antigen synthesis route	Reaction principle
Active site	Active group	Reaction method	Introduction group	Synthesis Method		
3	Hydroxyl	Acid anhydride reaction	carboxyl	MA		AFB1 is converted to AFB2a under the action of H ₂ SO ₄ . The 2-position hydroxyl of the active site is used to react with acid anhydride. The product is a half-ester compound AFB2a-HS. The carboxyl active group is introduced, and the coupling agent isobutyl chloroformate (IBC), AFB ₁ O and BSA synthesize AFB ₁ -BSA in the form of a single amide bond [11,12].
3	Aldehyde	Schiff	Aldehyde	SA		The condensation reaction of AFB1 under the action of H ₂ SO ₄ produces AFB2a with active sites of aldehyde groups, whose aldehyde groups can form unstable Schiff bases with the amino groups of BSA. Through the reduction of NaBH ₄ , the antigen AFB2a-BSA is synthesized [13,14].
3、4	Bifuran ring	Oxidation	Hydroxyl	EP		Using dichloromethane as solvent, the double bond of AFB1 bifuran ring is oxidized to form AFB1 epoxide, which reacts with the primary amine of BSA to form secondary amine, introduces a hydroxyl group on the epoxide, and couples with BSA in the form of monoamide into AFB1-BSA [15,16].
3、4	Bifuran ring	Glycolic acid	carboxyl	EED		The molecular structure of AFB1 contains an active site bifuran ring, which can react with glycolic acid to generate AFB1 enol ether derivative (AFB1-GA) with active carboxyl group, which is used to couple the carboxyl group with BSA to synthesize AFB1-BSA [17].

BGAFs artificial antigen identification

UV Scan

Dissolve AFB₁ with methanol, prepare 1 mg·mL⁻¹ AFB₁ solution; use volume ratio (v/v) 4:6 methanol PBS solution to dissolve BSA and AFB₁-BSA, prepare 1 mg·mL⁻¹ BSA and

AFB₁- BSA solution; UV scan at a wavelength of 200 ~ 500 nm, through the calculation formula $A = \epsilon CL$ (where A is the absorbance value, read by the instrument; ϵ is the molar extinction coefficient, which is a constant value; C is the solute concentration in the solution; L is Optical path, determined by the instru-

ment), calculate the molecular binding ratio of AFB₁ and BSA (Wang Y.N. et. al., 2014).

SDS-PAGE identification

The concentration of the concentrated gel and the separating gel are selected to be 5% and 12%, the voltage is 90 v and 60 v, the sample volume is 10 µL per well, and the protein content is 10 µg per well. The UV analyzer system software calculates AFB₁ and BSA the molecular binding ratio.

Preparation of AFB₁ pAb

The artificial antigens synthesized by 6 different methods were used to immunize New Zealand white rabbits. Each antigen was used to immunize 1 group, a total of 6 groups, 3 rabbits in each group. The immunization dose is calculated according to the amount of protein BSA in AFB₁-BSA, each is 100 µg, the volume is 1 mL, the back is injected subcutaneously at 4 to 6 points, a total of 5 immunizations, each interval is 3 to 4 weeks, after the fifth immunization for 2 weeks, blood was collected from the ear vein, the polyanterum was separated by centrifugation, and the polyanterum was purified by the saturated ammonium sulfate salting-out method to prepare AFB₁ pAb (Ju RH et. al., 2015).

Characteristic analysis of AFB₁ pAb

Determination of potency

Indirect ELISA (Zhao HH et. al., 2015).

Sensitivity identification

Indirect competitive ELISA (icELISA) measures the half inhibitory concentration (IC₅₀) of AFB₁ pAb on AFB₁ to determine sensitivity (Chen T et. al., 2014).

Specific identification

With AFB₁, AFB₂, AFG₁, and AFG₂ as inhibitors, the IC₅₀ of each inhibitor was determined by icELISA, and the percentage of the IC₅₀ of AFB₁ pAb to AFB₁ and the IC₅₀ of other inhibitors was used as the cross-reaction rate (CR%) (Zhang C et. al., 2016), the calculation method is $CR\% = IC_{50} \text{ of AFB}_1 \text{ pAb to AFB}_1 / IC_{50} \text{ of AFB}_1 \text{ pAb to other inhibitors}$.

Results

GAFs artificial antigen identification results

UV identification

The results are shown in Figure 2. In the range of UV200-500 nm, the characteristic peak of BSA is at 278 nm, and the characteristic peak of AFB₁ is at 363 nm. The artificial antigen AFB₁-BSA is synthesized by 6 methods including OAE, MOA, MA, SA, EP, EED. Both contain the characteristic peaks of BSA and AFB₁, indicating that the above 6 methods can synthesize artificial antigen AFB₁-BSA. The calculated results of the molecular binding ratio of BSA to AFB₁ (Liu, H.X. et. al., 2014) are shown in Table 2.




Fig.2 UV spectra of AFB₁-BSA

Table2

Molecular binding ratio of AFB₁-BSA prepared by six methods

Synthesis methods	Initial molar ratio of AFB ₁ to BSA	Molecular binding ratio of AFB ₁ -BSA	Usage ratio of AFB ₁
OAE	50:1	8.64:1	17.28
MOA	50:1	6.88:1	13.76
MA	50:1	10.78:1	21.56
SA	50:1	4.46:1	8.92
EP	50:1	6.38:1	12.76
EED	50:1	2.31:1	4.62

Note: Compared to the molecular weight of BSA and AFB₁, BSA is 66.446, AFB₁ is 312, BSA is much larger than AFB₁, so the utilization rate of BSA is 100% when the utilization

ratio is calculated.

SDS-PAGE identification

The results are shown in Figure 3. It can be seen that

the bands of the 6 artificial antigens AFB₁-BSA lag behind the bands of BSA, indicating that the molecular weight of AFB₁-

BSA is greater than that of BSA, and it can be determined that the synthesis of AFB₁-BSA is successful.




Fig.3 The SDS-PAGE photo of AFB₁-BSA

1. Maker; 2. BSA; 3. AFB₁-BSA(OAE); 4. AFB₁-BSA(MOA); 5. AFB₁-BSA(MA); 6. AFB₁-BSA(SA) ; 7. AFB₁-BSA(EP) ; 8. AFB₁-BSA(EED).

AFB₁ pAb characteristic analysis

Determination of potency

The results are shown in Figure 4. It can be seen that after the immunization, one rabbit with the highest indirect ELISA titer was selected for comparison and analysis in each

group. The indirect ELISA titer of the 6 immunized rabbits all reached 1: (1×10⁴). It can be seen that the 6 types The AFB₁-BSA synthesized by the method has good immunogenicity. The OAE group and MA group have the best immune effect, with a titer of 1: (1.28×10⁴).




Fig.4 The indirect ELISA titer curves of AFB₁ pAb

Sensitivity analysis

The results are shown in Figure 5. It can be seen that the icELISA inhibition curve of 6 immunized rabbits has a good

linear relationship. The OAE group has the best sensitivity, with an IC_{50} of $10.32 \mu\text{g}\cdot\text{kg}^{-1}$. The sensitivity of the other groups is inferior to that of the OAE group.




Fig.5 The sensitivity measurement of AFB₁ pAb to AFB₁ by icELISA

Table 3

The regression equation, R^2 and IC_{50} of AFB₁ pAb to AFB₁ by icELISA

group	Regression equation	R^2 value	IC_{50} ($\mu\text{g}\cdot\text{kg}^{-1}$)
OAE	$y = -32.171x + 82.621$	0.9916	10.32
MOA	$y = -31.822x + 99.59$	0.9943	36.18
MA	$y = -31.546x + 97.263$	0.9938	31.49
SA	$y = -32.875x + 92.292$	0.9966	19.36
EP	$y = -25.245x + 99.481$	0.9932	91.21

Specificity and broad-spectrum analysis

The results are shown in Table 4. It can be seen that the antibodies prepared by the six methods can recognize AFB₁ 100%, and the OAE method has the best specificity and broad-spectrum, with an IC_{50} of $10.32 \mu\text{g}\cdot\text{kg}^{-1}$ and a CR with AFB₂ of 86.46%; The CR with AFG₁ and AFG₂ were 44.13% and 14.72%, respectively. Antibodies prepared by other methods

have good specificity and can recognize AFB₁ 100%, but their sensitivity and broad-spectrum are not as good as those prepared by OAE method. The results show that the best antigen synthesis method for preparing antibodies against BGAFs with high sensitivity, strong specificity and good broad spectrum is the OAE method.

Table 4

The percent cross-reactivity of AFB₁ pAb with AFB₁, AFB₂, AFG₁, AFG₂

AF	AFB ₁ pAb(OAE)		AFB ₁ pAb(MOA)		AFB ₁ pAb(MA)		AFB ₁ pAb(SA)		AFB ₁ pAb(EP)		AFB ₁ pAb(EED)	
	IC_{50} ($\mu\text{g}\cdot\text{kg}^{-1}$)	(%) CR	IC_{50} ($\mu\text{g}\cdot\text{kg}^{-1}$)	(%) CR	IC_{50} ($\mu\text{g}\cdot\text{kg}^{-1}$)	(%) CR	IC_{50} ($\mu\text{g}\cdot\text{kg}^{-1}$)	(%) CR	IC_{50} ($\mu\text{g}\cdot\text{kg}^{-1}$)	(%) CR	IC_{50} ($\mu\text{g}\cdot\text{kg}^{-1}$)	(%) CR
AFB ₁	10.32	100	36.18	100	31.49	100	19.36	100	91.21	100	307.81	100
AFB ₂	13.72	75.21	46.03	78.61	46.85	67.22	23.83	81.26	144.14	63.28	590.13	52.16
AFG ₁	23.39	44.13	> 10^3	< 0.5	> 10^3	< 0.5	35.67	54.27	> 10^3	< 0.5	> 10^3	< 0.5
AFG ₂	70.11	14.72	> 10^3	< 0.5	> 10^3	< 0.5	79.41	24.38	> 10^3	< 0.5	> 10^3	< 0.5
AFM	63.08	16.36	741.39	4.88	615.04	5.12	541.74	3.68	> 10^3	< 0.5	> 10^3	< 0.5
AFM	716.67	1.44	> 10^3	< 0.5	> 10^3	< 0.5	> 10^3	< 0.5	> 10^3	< 0.5	> 10^3	< 0.5

Discussion and Conclusion

About the design of BGAFs antigen synthesis method

The molecular weights of AFB₁ and AFB₂ in BGAFs are 312.27 and 314.29, respectively. They belong to small molecule haptens and have no immunogenicity. According to the theory of hapten-carrier effect, only by combining with large-molecule protein carriers to form artificial antigens can they be specific for haptens. Therefore, the design of antigen synthesis methods is very important (Zeng, H. et. al., 2014). Since the selection of different active sites and the introduction of different linking arm

lengths will have a greater impact on the properties and structure of small molecules, which in turn will affect the quality of antibodies produced (Shi HY et. al., 2006). According to the molecular structure characteristics of BGAFs, this study selected the 1-position carbonyl group, 2-position active hydrogen, 3-position hydroxyl group and aldehyde group, and the difuran ring between 3-position and 4-position as the active groups. Through different chemical reaction methods, respectively introduce available carboxyl, hydroxyl, aminomethyl and other active groups to realize the coupling with carrier protein to synthesize

artificial antigens.

About the synthetic route of BGAFs artificial antigen

At present, the research on BGAFs artificial antigen synthesis method is still at the empirical level, and trial and error methods are mostly used. Although a variety of artificial antigen identification methods have been established, the immunogenicity of the artificial antigens prepared is ultimately through the effect of animal immunity. It was confirmed (Guo N. F. et. al., 2014). Based on a large number of relevant research literature, this article uses AFB₁ as the starting material for the reaction, and uses 6 methods such as OAE method, MOA method, MA method, SA method, EP method and EED method to synthesize artificial antigens, and through UV, SDS-PAGE for antigen identification and animal immunization for antibody characteristics analysis, the most ideal antigen synthesis method for the preparation of BGAFs antibody was selected by OAE method. Its advantages are that the reaction system is easy to construct, the reaction conditions are mild, the operation steps are simple, and the product yield is high. However, in terms of the advanced nature of the technical route adopted in this research, the research and application of molecular simulation technology, computer-aided technology, etc. Needs to be improved (Morita, I. 2017).

Analysis on the immune effect of BGAFs artificial antigen.

The purpose of this research is to screen out BGAFs artificial antigen synthesis methods, and lay the material and technical foundation for the preparation of high-quality BGAFs antibodies with high sensitivity, strong specificity and broad recognition spectrum. This requires that in the design of BGAFs antigen synthesis, on the one hand, it is necessary to consider the specificity and sensitivity of the antibody to AFB₁ to meet the detection technology requirements under the AFB₁ limit standard; On the other hand, it is necessary to consider the sensitivity and broad-spectrum of the antibody to AFB₂ to meet the technical requirements for detection under the BGAFs limit standard (Xie Hui et al. 2017) used MA method to synthesize AFB₁-BSA, and screened hybridoma cell 3B9 to obtain AFB₁ mab. The antibody specifically recognizes AFB₁ with a sensitivity of 1.04 $\mu\text{g}\cdot\text{kg}^{-1}$, CR of AFB₂, AFG₁, AFG₂, and AFM₁ are 2.2%, 33.9%, 1.8%, and 5.12%, respectively, which have no CR with AFM₂

and poor broad-spectrum. Xiao Zhi et al. used SA method to synthesize AFB₁-BSA, and screened hybridoma cell 3A12 to obtain AFB₁ mab. The antibody specifically recognizes AFB₁ with a sensitivity of 6.1 $\mu\text{g}\cdot\text{kg}^{-1}$, and is compatible with CR of AFB₂, AFG₁, AFG₂, and AMF₁. They are 7.8%, 20.2%, 0.6%, and 3.68%, respectively. It has no CR with AFM₂, and it also has the problem of poor broad-spectrum.

Conclusion

The results of 6 different antigen synthesis methods and the characteristics of the antibodies produced showed that the OAE method was the best, the produced AFB₁ pAb antibody titer was high, and the indirect ELISA titer reached 1: (1.28 $\times 10^4$); the sensitivity to AFB₁ was good, IC₅₀ is 10.32 $\mu\text{g}\cdot\text{kg}^{-1}$; it has strong specificity and can recognize AFB₁ 100%. The CR with AFB₂, AFG₁, AFG₂, AMF₁, and AFM₂ are 75.21%, 44.13%, 14.72%, 16.36% and 1.44%, respectively. The other five methods designed by this research have certain defects in varying degrees. Therefore, the author recommends that they should not be used except for research work.

In this study, based on the molecular structure characteristics of AFB₁ and the existing active sites, six BGAFs antigen synthesis methods were designed, through UV, SDS-PAGE identification and analysis of the characteristics of AFB₁ pAb produced by immunized animals, a high-titer, sensitive, specific, and broad-spectrum AFB₁ pAb was obtained. It shows that antigen synthesis design is the prerequisite for the preparation of high-quality antibodies, and the OAE method is an effective way to realize the preparation of high-quality antibodies for BGAFs, laying a material and technical foundation for the establishment of BGAFs immunoassay methods.

Author's contributions

All authors participated in this article design. Yanan WANG participated and performed writing and data collection. All authors read and approved the final manuscript. All authors contributed to the draft of the manuscript. All authors gave final approval for publication.

Conflict of interest Author does not report any financial or personal connections with other persons or organizations, which might negatively affect the contents of this publication and/or claim authorship rights to this publication.

References:

1. Sun, D.D.; Gu, X.; Li, J.G.; Yao, T.; Dong, Y.C. (2015). Quality evaluation of five commercial enzyme linked immunosorbent assay kits for detecting aflatoxin b1 in feedstuffs. *Asian-Australasian journal of animal sciences* 28, 691-696, doi:10.5713/ajas.14.0868.
2. Luo, X.; Li, K.; Xing, J.; Qi, L.; Yang, M.; Wang, R.; Wang, L.; Li, Y.; Chen, Z. (2018). In vivo toxicity assessment of aflatoxin B(1)-contaminated corn after ozone degradation. *Food Addit Contam Part A Chem Anal Control Expo Risk Assess* 35, 341-350, doi:10.1080/19440049.2017.1395518.
3. CHINA. National Food Safety Standard Limit of mycotoxins in food. (2017). Vol. GB 2761 —2017.
4. Gefen, T.; Vaya, J.; Khatib, S.; Rapoport, I.; Lupo, M.; Barnea, E.; Admon, A.; Heller, E.D.; Aizenshtein, E.; Pitcovski, J. (2015). The effect of haptens on protein-carrier immunogenicity. *Immunology* 144, 116-126, doi:10.1111/imm.12356.
5. Mongkon, W.; Sugita-Konishi, Y.; Chaisri, W.; Suriyasathaporn, W. (2017). Aflatoxin B1 Contamination of Dairy Feeds after Storage in Farm Practice in Northern Thailand. *Biocontrol Sci* 22, 41-45, doi:10.4265/bio.22.41.
6. Xiao LW, Xu X, Zhao W, et al. (2017). Study on the development and performance of a paper strip for rapid quantitative detection of aflatoxin B1 based on time-resolved fluorescent nano-spheres. *Grain Processing*. 42, 41-43.
7. Kolosova, A.Y.; Shim, W.B.; Yang, Z.Y.; Eremin, S.A.; Chung, D.H. (2006). Direct competitive ELISA based on a monoclonal antibody for detection of aflatoxin B1. Stabilization of ELISA kit components and application to grain samples. *Anal Bioanal Chem* 384, 286-294, doi:10.1007/s00216-005-0103-9.
8. Yu, Y.Y.; Chen, Y.Y.; Gao, X.; Liu, Y.Y.; Zhang, H.Y.; Wang, T.Y. (2018). Nanoparticle based bio-bar code technology for trace analysis of aflatoxin B1 in Chinese herbs. *Journal of food and drug analysis*. 26, 815-822, doi:10.1016/j.jfda.2017.11.003.

9. Wang YN, Wang XF, Niu LL, et al. (2016). Research progress in hapten molecule design and immunogen synthesis and antibody characteristics of aflatoxin B1. *Science and Technology of Food Industry*. 23, 367-376.
10. Zhou, Y.; Wu, J.; Yu, W.; Xu, Y.; Wang, P.; Xie, B.; Chen, F. (2007). Preparation for aflatoxin B(1)-cationized bovine serum albumin based on Mannich-type reaction. *J Immunol Methods*. 328, 79-88, doi:10.1016/j.jim.2007.08.009.
11. Rushing, B.R.; Selim, M.I. (2017). Structure and Oxidation of Pyrrole Adducts Formed between Aflatoxin B(2a) and Biological Amines. *Chem Res Toxicol*, 30, 1275-1285, doi:10.1021/acs.chemrestox.7b00002.
12. Kononenko, G.P.; Burkin, A.A.; Soboleva, N.A. (2002). [Comparative characteristics of immunoreagents based on aflatoxin B1 hemiacetals and sterigmatocystine]. *Prikl Biokhim Mikrobiol* 38, 571-577.
13. Wang YN, Wang XF, Niu LL, et al. (2017). Advance in immunoassay of total aflatoxins in food. *Science and Technology of Food Industry*. 13, 344-351.
14. Wang YN, Wang XF, Wang ZL. (2018). Study and application of detection methods of total aflatoxins in food. *Food and Fermentation Industries*. 44, 1, 285-290.
15. Cervino, C.; Knopp, D.; Weller, M.G.; Niessner, R. (2007). Novel aflatoxin derivatives and protein conjugates. *Molecules* 12, 641-653, doi:10.3390/12030641.
16. Wang YN, Wang SY, Zhang HT, et al. (2014). Establishment of Hybridoma Cell Lines Secreting Anti-Cadmium Ion Monoclonal Antibody and Identificat of Their Immunological Properties. *Acta Agriculturae Boreali-occidentalis Sinica.*, 11, 24-29.
17. Ju RH, Wen K, Duan LL, et al. (2015). Preparation and Characterization of Polyclonal Antibodies Against Acrylamide. *Journal of Nuclear Agricultural Sciences*. 29, 10, 1979-1984.
18. Zhao HH, Zhang ZR, Li XJ, et al. (2016). Prokaryotic Expression of Tartary Buckwheat Flavonol SynthaseFtFLS2 and Preparation of Its Polyclonal Antibody. *Journal of Nuclear Agricultural Sciences*. 30, 2, 240-245.
19. Chen T, Wang CQ, Li XF, et al. (2014). Determination of aflatoxin B1 in peanut under different storage conditions by indirect ELISA. *China Oils and Fats*. 9, 88-91.
20. Zhang C, Pan JR, Shuai RQ, et al. (2016). Research on Enzyme Linked Immunosorbent Assay for Multi-residues of Nitroimidazoles in Foods of Animal Origin. *Journal of Nuclear Agricultural Sciences*. 30, 2, 323-331.
21. Liu, H.X.; Yang, Y.X.; Ma, M.G.; Wang, X.M.; Du, X.Z. (2015). Self-assembled Gold Nanoparticles Coating for Solid-Phase Microextraction of Ultraviolet Filters in Environmental Water. *Chinese Journal of Analytical Chemistry* 43, 207-211.
22. Zeng, H.; Chen, J.; Zhang, C.; Huang, X.A.; Sun, Y.; Xu, Z.; Lei, H. (2016). Broad-Specificity Chemiluminescence Enzyme Immunoassay for (Fluoro)quinolones: Hapten Design and Molecular Modeling Study of Antibody Recognition. *Anal Chem* 88, 3909-3916, doi:10.1021/acs.analchem.6b00082..
23. Shi HY, Wang MH. (2008). Effect of Hapten Space Arm Length on Immune Recognition. *Chinese Journal of Pesticide Science*. 2, 172-177.
24. Guo NF, Yu JC, Ma FM, et al. (2014). Study of Comparison the Commonly Used Artificial Antigen Identification Methods. *Journal of Food Science and Biotechnology*. 33 5,517-521.
25. Morita, I.; Oyama, H.; Yasuo, M.; Matsuda, K.; Katagi, K.; Ito, A.; Tatsuda, H.; Tanaka, H.; Morimoto, S.; Kobayashi, N. (2017). Antibody Fragments for On-Site Testing of Cannabinoids Generated via in Vitro Affinity Maturation. *Biological & pharmaceutical bulletin*, 40, 174-181, doi:10.1248/bpb.b16-00669.
26. Hayashi, N.; Saegusa, J.; Uto, K.; Oyabu, C.; Saito, T.; Sato, I.; Kawano, S.; Kumagai, S. (2016). Evaluation of a Computer-Aided Microscope System and Its Anti-Nuclear Antibody Test Kit for Indirect Immunofluorescence Assay. *Rinsho Byori*, 64, 142-151.
27. Wang YN. (2020). Preparation of Broad-Specificity Antibodies and Development of Immuno-chromatographic Strip for Detection of Total Aflatoxins in Animal Food. *Henan Institute of Science and Technology*.
28. Xie H, Zhang X, Wang X, et al. (2015). Preparation of anti-aflatoxin B1 monoclonal antibodies and its use in an indirect competitive ELISA for aflatoxin B1. *Microbiology China*. 10, 2033-2040.
29. Xiao Z, Li PW, Zhuang Q, et al. (2011). Production and characteristics of specialised monoclonal antibodies against aflatoxin B1. *Chinese Journal of Oil Crop Sciences*. 33, 001, 66-70.

Янан Ванг, аспірант Сумський НАУ, (Суми, Україна), Коледж наук про тварин та ветеринарну медицину, Інститут науки і технологій Хенань, (Сінсян, Китай)

Ганна Фотіна, доктор ветеринарних наук, професор, Сумський НАУ, (Суми, Україна)

Синтез та підготовка антигенів для отримання специфічних і євритопічних антитіл проти B-group афлатоксинів

Метою цього дослідження було вироблення антитіл до афлатоксинів групи В (BGAF) із сильною специфічністю та хорошою євритопічністю. Дослідження проводили в лабораторії безпеки та якості продуктів тваринництва Сумського НАУ, факультету ветеринарної медицини, Суми, Україна та на базі Науково-технічного інституту Хенань, Сінсян, Китай. Відповідно до молекулярної структури та активного центру афлатоксину В1 (AFB1), штучний антиген BGAFs AFB1-BSA готували 6 ма методами, такими як метод активного ефіру оксиму (OAE), метилювання аміаку (MOA), змішаний ангідрид (MA), напівфабрикат ацеталь (SA) епоксид (EP) та похідне енолового ефіру (EED) та ідентифікували за допомогою УФ та SDS-PAGE.

Поліклональні антитіла проти AFB1 (AFB1 pAb) готували шляхом імунізації новозеландських кролів AFB1-BSA, а титри AFB1 pAb виявляли за допомогою непрямого ІФА, чутливість AFB1 pAb аналізували за допомогою непрямого конкурентного ІФА (icELISA), специфічність та євритопічність AFB1 pAb аналізували за допомогою тесту перехресної

реактивності (CR). Результати показали, що AFB1-BSA був успішно синтезований, і найкращим був метод активного ефіру оксиму (OAE) із 6 методів синтезу штучного антигену BGAF, а його відношення кон'югації AFB1 до BSA становило близько 8,46Å¹. Імунна ефективність методу OAE була найкращою, його рAb AFB1 мав високі титри 1: (1,28×10⁴) з використанням методу непрямого ІФА, чутливість з 50% концентрацією інгібування (IC₅₀) 10,32 мкг / л до AFB1 за допомогою ісELISA та високий CR до AFB2 75,21%, AFG1 44,13%, AFG2 14,72%, AFM1 16,36% та AFM2 1,44% відповідно. У цьому дослідженні були підготовлені рAbs AFB1 з високим титром, чутливістю, специфічністю та еуритопічністю, що заклало важливу та технічну основу для створення імунологічного аналізу BGAF.

Ключові слова: афлатоксини групи В, конструкція синтезу антигену, поліклональні антитіла, аналіз, характеристики.

Дата надходження до редакції: 20.11.2020 р.